

ELLIPTIC AND K -THEORETIC STABLE ENVELOPES AND NEWTON POLYTOPES

R. RIMÁNYI^{*}, V. TARASOV[°], A. VARCHENKO[◊]

^{*}[◊]*Department of Mathematics, University of North Carolina at Chapel Hill
Chapel Hill, NC 27599-3250, USA*

[◊]*Faculty of Mathematics and Mechanics, Lomonosov Moscow State University
Leninskiye Gory 1, 119991 Moscow GSP-1, Russia*

[°]*Department of Mathematical Sciences, Indiana University–Purdue University Indianapolis
402 North Blackford St, Indianapolis, IN 46202-3216, USA*

[°]*St. Petersburg Branch of Steklov Mathematical Institute
Fontanka 27, St. Petersburg, 191023, Russia*

ABSTRACT. In this paper we consider the cotangent bundles of partial flag varieties. We construct the K -theoretic stable envelopes for them and also define a version of the elliptic stable envelopes. We expect that our elliptic stable envelopes coincide with the elliptic stable envelopes defined by M. Aganagic and A. Okounkov. We give formulas for the K -theoretic stable envelopes and our elliptic stable envelopes. We show that the K -theoretic stable envelopes are suitable limits of our elliptic stable envelopes. That phenomenon was predicted by M. Aganagic and A. Okounkov. Our stable envelopes are constructed in terms of the elliptic and trigonometric weight functions which originally appeared in the theory of integral representations of solutions of qKZ equations twenty years ago. (More precisely, the elliptic weight functions had appeared earlier only for the \mathfrak{gl}_2 case.) We prove new properties of the trigonometric weight functions. Namely, we consider certain evaluations of the trigonometric weight functions, which are multivariable Laurent polynomials, and show that the Newton polytopes of the evaluations are embedded in the Newton polytopes of the corresponding diagonal evaluations. That property implies the fact that the trigonometric weight functions project to the K -theoretic stable envelopes.

CONTENTS

1. Introduction	2
2. Elliptic weight functions	5
2.1. Notation	5
2.2. Elliptic R -matrix	6
2.3. Index set \mathcal{I}_λ , variables t	6
2.4. Elliptic weight functions	6
2.5. Exchange properties	7
2.6. Transformation properties	10
2.7. Substitution properties of elliptic weight functions	12
2.8. Orthogonality	13

^{*}*E-mail:* rimanyi@email.unc.edu, supported in part by NSF grant DMS-1200685

[°]*E-mail:* vtarasov@iupui.edu, vt@pdmi.ras.ru, supported in part by Simons Foundation grant 430235

[◊]*E-mail:* anv@email.unc.edu, supported in part by NSF grants DMS-1362924, DMS-1665239

3.	Trigonometric weight functions	14
3.1.	Alcoves	14
3.2.	Elliptic versus trigonometric weight functions: the $q \rightarrow 0$ limit	15
3.3.	Recursion properties	16
3.4.	Substitution properties of trigonometric weight functions	16
3.5.	Newton polytope properties	17
4.	Partial flag varieties	20
4.1.	Definitions	20
4.2.	Schubert cells, conormal bundles	20
4.3.	Torus action	21
4.4.	Equivariant K -theory of $T^*\mathcal{F}_\lambda$	21
4.5.	Equivariant localization – moment map description	21
4.6.	Normal Euler classes of conormal spaces at torus fixed points	22
5.	K -theoretic stable envelopes	23
5.1.	Definition of K -theoretic stable envelopes	23
5.2.	Weight functions project to stable envelope classes	24
5.3.	Note on R -matrices	26
6.	Line bundles and quadratic forms	27
6.1.	Line bundles on \mathbb{C}^p	27
6.2.	Theta functions as sections of line bundles	28
7.	Equivariant elliptic cohomology of $T^*\mathcal{F}_\lambda$	29
7.1.	Tautological bundles and Chern classes	29
7.2.	Equivariant localization – moment map description	30
7.3.	Cotangent bundle and the dynamical parameters	30
7.4.	Admissible line bundles, and their sections	31
7.5.	T -equivariant elliptic cohomology classes	31
7.6.	Weight functions project to equivariant elliptic cohomology classes	32
7.7.	Elliptic stable envelope for cotangent bundles of partial flag varieties	32
7.8.	Axiomatic definition of the stable envelope	33
8.	Appendix: Comparison of Bethe algebras	35
	References	36

1. INTRODUCTION

Malik and Okounkov started in [MO1] a program to relate the quantum torus equivariant generalized cohomology (cohomology, K -theory, elliptic cohomology) of Nakajima varieties and representation theory of quantum groups. A central role in that program is played by the stable envelopes, which are maps from the equivariant generalized cohomology of the fixed point set of the torus action to the equivariant generalized cohomology of the variety. Stable envelopes depend on the choice of a chamber (a connected component of the complement of an arrangement of real hyperplanes) and stable envelopes of different chambers are related by R -matrices of the corresponding quantum group.

The basic example of a Nakajima variety is the cotangent bundle of a variety of partial flags in \mathbb{C}^n . Such a variety \mathcal{F}_λ is labeled by $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_N) \in \mathbb{Z}_{\geq 0}^N$, $|\boldsymbol{\lambda}| = \lambda_1 + \dots + \lambda_N = n$ and consists of flags

$$F_1 \subset F_2 \subset \dots \subset F_N = \mathbb{C}^n,$$

with $\dim F_i/F_{i-1} = \lambda_i$, $i = 1, \dots, N$. The torus is $T = (\mathbb{C}^\times)^n \times \mathbb{C}^\times$, with $(\mathbb{C}^\times)^n$ acting by diagonal matrices on \mathbb{C}^n and \mathbb{C}^\times acting by multiplication on the cotangent spaces. In this case the stable envelope map can be thought of as a linear map Stab_σ from $(\mathbb{C}^N)^{\otimes n}$ to the T -equivariant generalized cohomology of the disjoint union $\sqcup_{\lambda \in \mathbb{Z}_{\geq 0}^N, |\lambda|=n} T^* \mathcal{F}_\lambda$ of cotangent bundles. The stable envelope map depends on an element σ of the symmetric group S_n .

Let v_1, \dots, v_N be the standard basis of \mathbb{C}^N . The standard basis of $(\mathbb{C}^N)^{\otimes n}$ consists of vectors v_I labeled by partitions $I = (I_1, \dots, I_N)$ of $\{1, \dots, n\}$, $v_I = v_{i_1} \otimes \dots \otimes v_{i_n}$, where $i_a = j$ if $a \in I_j$. Let \mathcal{I}_λ be the set of all partitions I such that $|I_j| = \lambda_j$, $j = 1, \dots, N$. Then Stab_σ is the same as a collection of cohomology classes $(\kappa_{\sigma, I})_{I \in \mathcal{I}_\lambda}$ of $T^* \mathcal{F}_\lambda$ for all $\lambda \in \mathbb{Z}_{\geq 0}^N$, $|\lambda| = n$. The classes must satisfy some remarkable defining relations.

The existence and uniqueness of the stable envelopes for the T -equivariant cohomology of Nakajima varieties was established in [MO1]. The special case of the partial flag varieties see in [GRTV, RTV1], where the formulas for the stable envelopes were given. The K -theoretic stable envelopes were defined by Maulik and Okounkov in the paper [MO2], which is in preparation, see this definition in [O, OS] and in Section 5.1 below. According to [O, OS], the paper [MO2] will also contain the theorem of existence and uniqueness of the K -theoretic stable envelopes for Nakajima varieties. The proof of the uniqueness of the K -theoretic stable envelopes is also given in [O]. One of the main results of this paper is the proof of the existence of the K -theoretic stable envelopes for cotangent bundles of partial flag varieties and formulas for them, see Theorem 5.1.

The definition of elliptic stable envelopes for Nakajima varieties was sketched by Aganagic and Okounkov in [AO]. In [FRV] a version of the elliptic stable envelopes was defined for the cotangent bundles of Grassmannians. The second main result of this paper is the definition of a version of the elliptic stable envelopes for cotangent bundles of partial flag varieties in the spirit of [FRV] and an axiomatic description of them in Theorem 7.3. That definition coincides with the definition in [FRV] for the cotangent bundles of Grassmannians. We expect that the stable envelopes in [FRV] and this paper coincide with the corresponding elliptic stable envelopes in [AO].

Our formulas for different versions of stable envelopes are given in terms of the *weight functions*, which originally appeared in the theory of integral representations for solutions of different versions of qKZ equations and associated Bethe ansatz, see [SV, V, TV1, TV2, TV3, TV4, FTV1, FTV2, MTV]. The integral representation for a solution $I(z_1, \dots, z_n, y)$ of the Yangian $Y_y(\mathfrak{gl}_N)$ qKZ equation with values in $(\mathbb{C}^N)^{\otimes n}$ has the form

$$I(z_1, \dots, z_n, y) = \sum_I \left(\int \Phi(\mathbf{t}, z_1, \dots, z_n, y) W_I(\mathbf{t}, z_1, \dots, z_n, y) d\mathbf{t} \right) v_I,$$

where $\Phi(\mathbf{t}, \mathbf{z}, y)$ is some scalar *master function* and $W_I(\mathbf{t}, \mathbf{z}, y)$ are the weight functions, y is deformation parameter of the Yangian, $\mathbf{t} = (t_i^{(k)})$ are the integration variables. To obtain a T -equivariant cohomological stable envelope we identify the variables z_1, \dots, z_n, y with the equivariant parameters of the torus $T = (\mathbb{C}^\times)^n \times \mathbb{C}^\times$ and the variables \mathbf{t} with equivariant Chern roots of the tautological vector bundles on $T^* \mathcal{F}_\lambda$. To construct the K -theoretic or elliptic stable envelopes we take the weight functions appearing in other versions of the qKZ equations.

Initially the weight functions were invented to construct the integral representations for solutions of the qKZ equations. The main requirement on them was the condition that the weight functions $(W_{\sigma,I}(\mathbf{t}, \mathbf{z}, y))_{I \in \mathcal{I}_\lambda}$ and $(W_{\sigma',I}(\mathbf{t}, \mathbf{z}, y))_{I \in \mathcal{I}_\lambda}$ labeled by $\sigma, \sigma' \in S_n$ must be related by the R -matrices of the corresponding quantum group, which permute the factors in the tensor product of n evaluation representations. No stable envelope properties were expected from them. Only now, 20 years later, when the stable envelopes were introduced, the new stable envelope properties of the weight functions are becoming uncovered.

In this paper we study the \mathfrak{gl}_N elliptic and trigonometric weight functions. For $N = 2$, the elliptic weight functions were introduced in [FTV1, FTV2]. For $N > 2$, the \mathfrak{gl}_N elliptic weight functions did not appear previously in the literature. The \mathfrak{gl}_N trigonometric weight functions of this paper are modifications of the trigonometric weight functions considered in [TV1, TV4].

In this paper we consider, in particular, the trigonometric weight functions denoted by $(\tilde{W}_{\sigma,I}^\Delta(\mathbf{t}, \mathbf{z}, h))_{I \in \mathcal{I}_\lambda}$ and their evaluations $(\tilde{W}_{\sigma,I}^\Delta(\mathbf{z}_J, \mathbf{z}, h))_{I \in \mathcal{I}_\lambda}$ labeled by partitions $J \in \mathcal{I}_\lambda$. Each of these evaluations is a Laurent polynomial in z_1, \dots, z_n . We show that the diagonal evaluation $\tilde{W}_{\sigma,J}^\Delta(\mathbf{z}_J, \mathbf{z}, h)$ equals some explicit product of binomials and the Newton polytope of any off-diagonal evaluation $\tilde{W}_{\sigma,I}^\Delta(\mathbf{z}_J, \mathbf{z}, h)$ with $I \neq J$ can be parallelly moved inside the Newton polytope of the corresponding diagonal evaluation $\tilde{W}_{\sigma,J}^\Delta(\mathbf{z}_J, \mathbf{z}, h)$, see Theorem 3.9. This is the stable envelope property of the trigonometric weight functions. Theorem 3.9 is our third main result.

We emphasize that the weight functions are not the same object as the stable envelopes. The weight functions are polynomials of certain variables, while the stable envelopes are projections of the weight functions (usually divided by some nontrivial factors) to the corresponding equivariant cohomology algebras with relations.

Our fourth main result is the relation between our elliptic stable envelopes, introduced in this paper, and the trigonometric stable envelopes, see Theorem 3.1. That relation between the elliptic and K -theoretic stable envelopes was predicted in [AO, Proposition 3.5], see a remark after Theorem 3.1. The elliptic weight functions are defined as symmetrizations of alternating products of theta functions of one variable. When the modular parameter and the argument of the theta function tend to zero in a special way the theta function turns into a binomial. The corresponding limit of an elliptic weight function turns into a trigonometric weight function. In that way the trigonometric stable envelopes can be recovered from the elliptic stable envelopes.

The exposition is the following. In Section 2 we introduce the elliptic weight functions and study their properties. Theorem 2.2 shows that different elliptic weight functions are related by the \mathfrak{gl}_N elliptic dynamical R -matrices. Theorem 2.9 gives the orthogonality relations for the elliptic weight functions, similarly to the orthogonality relations in [TV2, TV3, RTV1, RTV2, FRV]. In Section 3 we introduce the trigonometric weight functions. Theorem 3.2 describes the R -matrix properties of the trigonometric weight functions. In Section 3 we formulate and prove Theorem 3.9 on Newton polytopes.

In Section 4 we discuss elementary facts on partial flag varieties. In Section 5 the K -theoretic stable envelopes of cotangent bundles of partial flag varieties are constructed. In Section 6 we remind elementary facts on line bundles on powers of an elliptic curve. In

Section 7 we define our elliptic stable envelopes. At the end of the paper we comment on the definitions of stable envelopes in cohomology, K -theory, and elliptic cohomology.

This paper was inspired by papers and oral presentations by Andrei Okounkov and his coauthors. Our goal was to understand the K -theoretic and elliptic stable envelopes and their relations with weight functions. The authors thank G. Felder for useful discussions and the referee for helpful suggestions. The authors thank H. Konno who pointed out a mistake in the earlier version of Theorem 2.9, in which the shift h^λ had been missing, see that shift in our earlier papers [RV1, FRV]. The third author thanks the Max Planck Institute for Mathematics and Hausdorff Institute for Mathematics in Bonn for hospitality.

Remark on notation. Weight functions appear in the literature in three flavors: rational, trigonometric, and elliptic. Their natural notation would be $W_{\sigma,I}^{rat}$, $W_{\sigma,I,\Delta}^{trig}$, $W_{\sigma,I}^{ell}$. We will use the latter for the elliptic version. For the trigonometric one, we will write $W_{\sigma,I}^\Delta$ to keep the notation simpler. There will be no rational weight functions in this paper.

2. ELLIPTIC WEIGHT FUNCTIONS

2.1. **Notation.** Let τ be a complex number with positive imaginary part and let $q = e^{2\pi i\tau}$. We will use the complex variables x and u satisfying $x = e^{2\pi iu}$. We set $q^{1/2} = e^{\pi i\tau}$ and $x^{1/2} = e^{\pi iu}$. Define the theta function

$$(2.1) \quad \vartheta(x) = (x^{1/2} - x^{-1/2}) \phi(qx) \phi(q/x), \quad \phi(x) = \prod_{s=0}^{\infty} (1 - q^s x),$$

cf. [AO, (67)]. Then

$$(2.2) \quad \frac{\vartheta(qx)}{\vartheta(x)} = -\frac{1}{q^{1/2}x}, \quad \vartheta(1/x) = -\vartheta(x),$$

Let

$$(2.3) \quad \theta(u) = \theta(u, \tau) = \vartheta(e^{2\pi iu}, e^{2\pi i\tau})$$

be obtained from $\vartheta(x, q)$ by the substitution $q = e^{2\pi i\tau}$, $x = e^{2\pi iu}$. Then

$$(2.4) \quad \theta(u+1) = -\theta(u), \quad \theta(u+\tau) = -e^{-\pi i\tau - 2\pi iu} \theta(u).$$

Let $q \rightarrow 0$, x, a, b are fixed, $0 < \operatorname{Re} \varepsilon < 1$, and $m \in \mathbb{Z}$, then

$$(2.5) \quad \vartheta(x) \rightarrow x^{1/2} - x^{-1/2}, \quad \frac{\vartheta(aq^{m+\varepsilon})}{\vartheta(bq^{m+\varepsilon})} \rightarrow (a/b)^{-m-1/2}.$$

The following identities holds for $\vartheta(x)$:

$$(2.6) \quad \begin{aligned} \vartheta(\alpha y_1/x) \vartheta(h y_2/x) \vartheta(h y_1/y_2) \vartheta(\alpha) = \\ = \vartheta(\alpha h y_1/x) \vartheta(y_2/x) \vartheta(y_1/y_2) \vartheta(\alpha/h) + \vartheta(h y_1/x) \vartheta(\alpha y_2/x) \vartheta(h) \vartheta(\alpha y_1/y_2), \end{aligned}$$

$$\begin{aligned}
(2.7) \quad & \vartheta(\alpha_1 \alpha_2 h y_1 / x_1) \vartheta(y_2 / x_1) \vartheta(h y_1 / x_2) \vartheta(\alpha_2 h y_2 / x_2) \vartheta(h x_2 / x_1) \vartheta(y_1 / y_2) \vartheta(\alpha_1) - \\
& - \vartheta(h y_1 / x_1) \vartheta(\alpha_2 h y_2 / x_1) \vartheta(\alpha_1 \alpha_2 h y_1 / x_2) \vartheta(y_2 / x_2) \vartheta(\alpha_1 x_2 / x_1) \vartheta(y_1 / y_2) \vartheta(h) = \\
& = \vartheta(\alpha_1 \alpha_2 h y_1 / x_1) \vartheta(h y_2 / x_1) \vartheta(y_1 / x_2) \vartheta(\alpha_2 h y_2 / x_2) \vartheta(x_2 / x_1) \vartheta(h y_1 / y_2) \vartheta(\alpha_1) - \\
& - \vartheta(h y_1 / x_1) \vartheta(\alpha_1 \alpha_2 h y_2 / x_1) \vartheta(\alpha_2 h y_1 / x_2) \vartheta(y_2 / x_2) \vartheta(x_2 / x_1) \vartheta(\alpha_1 y_1 / y_2) \vartheta(h).
\end{aligned}$$

2.2. Elliptic R -matrix. Given a positive integer N , let $\boldsymbol{\mu} = (\mu_1, \dots, \mu_N)$. Nonzero entries of Felder's dynamical R -matrix ${}^F\mathcal{R}(x, \boldsymbol{\mu})$ are

$${}^F\mathcal{R}_{jj}^{jj}(x, \boldsymbol{\mu}) = 1, \quad {}^F\mathcal{R}_{jk}^{jk}(x, \boldsymbol{\mu}) = \frac{\vartheta(x) \vartheta(h \mu_j / \mu_k)}{\vartheta(xh) \vartheta(\mu_j / \mu_k)}, \quad {}^F\mathcal{R}_{kj}^{jk}(x, \boldsymbol{\mu}) = \frac{\vartheta(x \mu_j / \mu_k) \vartheta(h)}{\vartheta(xh) \vartheta(\mu_j / \mu_k)},$$

where $j \neq k$, see [F], where $h = e^{-2\pi i \gamma}$ and $\mu_j = e^{-2\pi i \lambda_j}$, $j = 1, \dots, N$. The R -matrix $\mathcal{R}(x, \boldsymbol{\mu})$

$$(2.8) \quad \mathcal{R}_{jj}^{jj}(x, \boldsymbol{\mu}) = 1, \quad \mathcal{R}_{kj}^{jk}(x, \boldsymbol{\mu}) = \frac{\vartheta(x \mu_j / \mu_k) \vartheta(h)}{\vartheta(xh) \vartheta(\mu_j / \mu_k)}, \quad j \neq k,$$

$$\mathcal{R}_{jk}^{jk}(x, \boldsymbol{\mu}) = \frac{\vartheta(x) \vartheta(h \mu_j / \mu_k) \vartheta(h \mu_k / \mu_j)}{\vartheta(xh) \vartheta(\mu_j / \mu_k) \vartheta(\mu_k / \mu_j)}, \quad \mathcal{R}_{kj}^{kj}(x, \boldsymbol{\mu}) = \frac{\vartheta(x)}{\vartheta(xh)}, \quad j < k,$$

is similar (in the case $N = 2$) to the R -matrix in [AO, formula (66)] under the identification $x = u$, $h = 1/\hbar$, $\mu_1/\mu_2 = z$.

2.3. Index set \mathcal{I}_λ , variables \mathbf{t} . We will use the following notations throughout the paper. Let $n, N \in \mathbb{Z}_{\geq 0}$ and let $\boldsymbol{\lambda} \in \mathbb{Z}_{\geq 0}^N$ be such that $\sum_{k=1}^N \lambda_k = n$. Set $\lambda^{(k)} = \lambda_1 + \dots + \lambda_k$ for $k = 0, \dots, N$, and $\lambda^{\{1\}} = \lambda^{(1)} + \dots + \lambda^{(N-1)}$.

The set of partitions $I = (I_1, \dots, I_N)$ of $\{1, \dots, n\}$ with $|I_k| = \lambda_k$ is denoted by \mathcal{I}_λ . For $I \in \mathcal{I}_\lambda$ we will use the notation $\bigcup_{a=1}^k I_a = \{i_1^{(k)} < \dots < i_{\lambda^{(k)}}^{(k)}\}$.

Consider variables $t_a^{(k)}$ for $k = 1, \dots, N$, $a = 1, \dots, \lambda^{(k)}$, where $t_a^{(N)} = z_a$, $a = 1, \dots, n$. Denote $t^{(j)} = (t_k^{(j)})_{k \leq \lambda^{(j)}}$ and $\mathbf{t} = (t^{(1)}, \dots, t^{(N-1)})$.

2.4. Elliptic weight functions. For $I \in \mathcal{I}_\lambda$ define the *elliptic weight function*

$$(2.9) \quad W_I^{\text{ell}}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu}) = (\vartheta(h))^{\lambda^{\{1\}}} \text{Sym}_{t^{(1)}} \dots \text{Sym}_{t^{(N-1)}} U_I(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu}),$$

where $\text{Sym}_{t^{(k)}}$ is the symmetrization with respect to the variables $t_1^{(k)}, \dots, t_{\lambda^{(k)}}^{(k)}$,

$$\text{Sym}_{t^{(k)}} f(t_1^{(k)}, \dots, t_{\lambda^{(k)}}^{(k)}) = \sum_{\sigma \in S_{\lambda^{(k)}}} f(t_{\sigma(1)}^{(k)}, \dots, t_{\sigma(\lambda^{(k)})}^{(k)}),$$

$$(2.10) \quad U_I^{\text{ell}}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu}) = \prod_{k=1}^{N-1} \prod_{a=1}^{\lambda^{(k)}} \left(\prod_{c=1}^{\lambda^{(k+1)}} \psi_{I,k,a,c}^{\text{ell}}(t_c^{(k+1)} / t_a^{(k)}) \prod_{b=a+1}^{\lambda^{(k)}} \frac{\vartheta(ht_b^{(k)} / t_a^{(k)})}{\vartheta(t_b^{(k)} / t_a^{(k)})} \right),$$

$$(2.11) \quad \psi_{I,k,a,c}^{\text{ell}}(x) = \begin{cases} \vartheta(hx), & \text{if } i_c^{(k+1)} < i_a^{(k)}, \\ \frac{\vartheta(xh^{1+p_{I,j(I,k,a)}(i_a^{(k)})-p_{I,k+1}(i_a^{(k)})}\mu_{k+1}/\mu_{j(I,k,a)})}{\vartheta(h^{1+p_{I,j(I,k,a)}(i_a^{(k)})-p_{I,k+1}(i_a^{(k)})}\mu_{k+1}/\mu_{j(I,k,a)})}, & \text{if } i_c^{(k+1)} = i_a^{(k)}, \\ \vartheta(x), & \text{if } i_c^{(k+1)} > i_a^{(k)}, \end{cases}$$

where $j(I, k, a) \in \{1, \dots, N\}$ is such that $i_a^{(k)} \in I_{j(I,k,a)}$, and

$$(2.12) \quad p_{I,j}(m) = |I_j \cap \{1, \dots, m-1\}|, \quad j = 1, \dots, N.$$

Denote by $\psi_I(h, \boldsymbol{\mu})$ the product of all denominators appearing in (2.10) and depending on $h, \boldsymbol{\mu}$ only,

$$(2.13) \quad \psi_I(h, \boldsymbol{\mu}) = \prod_{k=1}^{N-1} \prod_{k=1}^{\lambda^{(k)}} \vartheta(h^{1+p_{I,j(I,k,a)}(i_a^{(k)})-p_{I,k+1}(i_a^{(k)})}\mu_{k+1}/\mu_{j(I,k,a)}).$$

Remark. Notice that the weight functions are regular at the diagonals $t_b^{(k)} = t_a^{(k)}$ despite the appearance of the denominators $\vartheta(t_b^{(k)}/t_a^{(k)})$ in (2.10).

For $\sigma \in S_n$ and $I \in \mathcal{I}_\lambda$, define the elliptic weight function

$$(2.14) \quad W_{\sigma,I}^{\text{ell}}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu}) = W_{\sigma^{-1}(I)}^{\text{ell}}(\mathbf{t}, z_{\sigma(1)}, \dots, z_{\sigma(n)}, h, \boldsymbol{\mu}),$$

where $\sigma^{-1}(I) = (\sigma^{-1}(I_1), \dots, \sigma^{-1}(I_N))$. Hence, $W_I^{\text{ell}} = W_{\text{id},I}^{\text{ell}}$.

Example. For $N = 2, n = 2, \lambda = (1, 1), s = (1, 2) \in S_2, t_1^{(1)} = t$, we have

$$\begin{aligned} W_{\text{id},(\{1\},\{2\})}^{\text{ell}} &= \vartheta(h)\vartheta(z_2/t) \frac{\vartheta(hz_1/t \cdot \mu_2/\mu_1)}{\vartheta(h\mu_2/\mu_1)}, \\ W_{\text{id},(\{2\},\{1\})}^{\text{ell}} &= \vartheta(h)\vartheta(hz_1/t) \frac{\vartheta(z_2/t \cdot \mu_2/\mu_1)}{\vartheta(\mu_2/\mu_1)}, \\ W_{s,(\{1\},\{2\})}^{\text{ell}} &= \vartheta(h)\vartheta(hz_2/t) \frac{\vartheta(z_1/t \cdot \mu_2/\mu_1)}{\vartheta(\mu_2/\mu_1)}, \\ W_{s,(\{2\},\{1\})}^{\text{ell}} &= \vartheta(h)\vartheta(z_1/t) \frac{\vartheta(hz_2/t \cdot \mu_2/\mu_1)}{\vartheta(h\mu_2/\mu_1)}. \end{aligned}$$

2.5. Exchange properties. Let

$$(2.15) \quad \beta(x_1, x_2, y_1, y_2, \alpha) = \text{Sym}_{x_1, x_2} \vartheta(\alpha y_1/x_1) \vartheta(y_2/x_1) \vartheta(hy_1/x_2) \vartheta(\alpha hy_2/x_2) \frac{\vartheta(hx_2/x_1)}{\vartheta(x_2/x_1)}.$$

Lemma 2.1. $\beta(x_1, x_2, y_1, y_2, \alpha)$ is symmetric in y_1, y_2 .

Proof. Identity (2.7) for $\alpha_1 = 1/h, \alpha_2 = \alpha$ is equivalent to

$$(2.16) \quad \beta(x_1, x_2, y_1, y_2, \alpha) = \beta(\alpha hy_1, \alpha hy_2, x_1, x_2, \alpha),$$

which proves the claim. \square

Let $s_{i,j} \in S_n$ denote the transposition of i and j . Set $\boldsymbol{\mu}_I^{(i)} = (h^{-p_{I,1}(i)}\mu_1, \dots, h^{-p_{I,N}(i)}\mu_N)$.

Theorem 2.2. *Let $\sigma \in S_n$ be such that $\sigma(i) \in I_a$ and $\sigma(i+1) \in I_b$. Then*

$$(2.17) \quad W_{\sigma s_{i,i+1}, I}^{\text{ell}} = W_{\sigma, I}^{\text{ell}}$$

for $a = b$, and

$$(2.18) \quad W_{\sigma s_{i,i+1}, I}^{\text{ell}} = \mathcal{R}_{ab}^{ab}(z_{\sigma(i)}/z_{\sigma(i+1)}, \boldsymbol{\mu}_{\sigma^{-1}(I)}^{(i)}) W_{\sigma, I}^{\text{ell}} + \mathcal{R}_{ab}^{ba}(z_{\sigma(i)}/z_{\sigma(i+1)}, \boldsymbol{\mu}_{\sigma^{-1}(I)}^{(i)}) W_{\sigma, s_{\sigma(i), \sigma(i+1)}(I)}^{\text{ell}}$$

for $a \neq b$.

Proof. By formula (2.14), it suffices to prove the statement for σ being the identity permutation. In that case, $i \in I_a$, $i+1 \in I_b$, formula (2.17) for $a = b$ reads

$$(2.19) \quad W_I^{\text{ell}}(\mathbf{t}, \mathbf{z}^{(i)}, h, \boldsymbol{\mu}) = W_I^{\text{ell}}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu}),$$

where $\mathbf{z}^{(i)} = (z_1, \dots, z_{i-1}, z_{i+1}, z_i, z_{i+2}, \dots, z_n)$, and formula (2.18) for $a \neq b$ reads

$$(2.20) \quad W_{s_{i,i+1}(I)}^{\text{ell}}(\mathbf{t}, \mathbf{z}^{(i)}, h, \boldsymbol{\mu}) = \mathcal{R}_{ab}^{ab}(z_i/z_{i+1}, \boldsymbol{\mu}_I^{(i)}) W_I^{\text{ell}}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu}) + \\ + \mathcal{R}_{ab}^{ba}(z_i/z_{i+1}, \boldsymbol{\mu}_I^{(i)}) W_{s_{i,i+1}(I)}^{\text{ell}}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu}).$$

Furthermore, formula (2.10) implies that it suffices to consider only the case $n = 2$, $i = 1$. By formula (2.10), the products $U_{s_{i,i+1}(I)}^{\text{ell}}(\mathbf{t}, \mathbf{z}^{(i)}, h, \boldsymbol{\mu})$, $U_I^{\text{ell}}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu})$, and $U_{s_{i,i+1}(I)}^{\text{ell}}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu})$ have many common factors, being different in the part that reproduces the case $n = 2$, $i = 1$ up to a change of notation. Then formula (2.20) in general can be obtained by taking formula (2.20) for $n = 2$, $i = 1$ in appropriate variables, multiplying it by the common factors of the products $U_{s_{i,i+1}(I)}^{\text{ell}}(\mathbf{t}, \mathbf{z}^{(i)}, h, \boldsymbol{\mu})$, $U_I^{\text{ell}}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu})$, $U_{s_{i,i+1}(I)}^{\text{ell}}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu})$, and subsequent symmetrization in $\mathbf{t}^{(j)}$ variables for each $j = 1, \dots, N-1$.

Assume $n = 2$, $i = 1$. To simplify the notation, we write $s = s_{1,2}$. The proof uses the recursive structure of the weight functions. Set

$$\tilde{U}_I^{(k)}(\mathbf{t}^{(k)}, \mathbf{t}^{(k+1)}) = \prod_{a=1}^{\lambda^{(k)}} \left(\prod_{c=1}^{\lambda^{(k+1)}} \psi_{k,a,c}^{\text{ell}}(t_c^{(k+1)}/t_a^{(k)}) \prod_{b=a+1}^{\lambda^{(k)}} \frac{\vartheta(ht_b^{(k)}/t_a^{(k)})}{\vartheta(t_b^{(k)}/t_a^{(k)})} \right),$$

so that $U_I^{\text{ell}}(\mathbf{t}, \mathbf{z}) = \prod_{k=1}^{N-1} \tilde{U}_I^{(k)}(\mathbf{t}^{(k)}, \mathbf{t}^{(k+1)})$, see (2.10), (2.11). Define

$$(2.21) \quad W_I^{(k)}(\mathbf{t}^{(k-1)}, \mathbf{t}^{(k)}) = \text{Sym}_{\mathbf{t}^{(1)}} \dots \text{Sym}_{\mathbf{t}^{(k-1)}} (\tilde{U}_I^{(1)}(\mathbf{t}^{(1)}, \mathbf{t}^{(2)}) \dots \tilde{U}_I^{(k-1)}(\mathbf{t}^{(k-1)}, \mathbf{t}^{(k)}))$$

where $\mathbf{t}^{(j)} = (t^{(1)}, \dots, t^{(j)})$. In particular,

$$(2.22) \quad W_I^{\text{ell}}(\mathbf{t}, \mathbf{z}) = (\vartheta(h))^{\lambda^{\{1\}}} W_I^{(N)}(\mathbf{t}, \mathbf{z}).$$

Formula (2.21) yields

$$(2.23) \quad W_I^{(k+1)}(\mathbf{t}^{(k)}, \mathbf{t}^{(k+1)}) = \text{Sym}_{\mathbf{t}^{(k)}} (W_I^{(k)}(\mathbf{t}^{(k-1)}, \mathbf{t}^{(k)}) \tilde{U}_I^{(k)}(\mathbf{t}^{(k)}, \mathbf{t}^{(k+1)})).$$

Let $a = b$. Then $I_a = \{1, 2\}$, $I_c = \emptyset$ for $c \neq a$, $\tilde{U}_I^{(k)} = 1$ for $k < a$, and

$$\tilde{U}_I^{(k)} = \zeta(t_1^{(k)}, t_2^{(k)}, t_1^{(k+1)}, t_2^{(k+1)}, 1/h, h\mu_{k+1}/\mu_a)$$

for $k \geq a$, where

$$(2.24) \quad \zeta(x_1, x_2, y_1, y_2, \alpha_1, \alpha_2) = \vartheta(\alpha_1 \alpha_2 h y_1 / x_1) \vartheta(y_2 / x_1) \vartheta(h y_1 / x_2) \vartheta(\alpha_2 h y_2 / x_2) \frac{\vartheta(h x_2 / x_1)}{\vartheta(x_2 / x_1)},$$

cf. (2.15). Therefore, by formula (2.23) and Lemma 2.1,

$$W_I^{(N)}(\mathbf{t}, z_1, z_2) = \prod_{k=a}^{N-1} \beta(t_1^{(k)}, t_2^{(k)}, t_1^{(k+1)}, t_2^{(k+1)}, h \mu_{k+1} / \mu_1) = W_I^{(N)}(\mathbf{t}, z_2, z_1).$$

By formulae (2.14) and (2.22), we have $W_{s,I}^{\text{ell}}(\mathbf{t}, z_1, z_2) = W_I^{\text{ell}}(\mathbf{t}, z_2, z_1) = W_I^{\text{ell}}(\mathbf{t}, z_1, z_2)$, which proves equality (2.19).

Let $a \neq b$. Then $I_a = \{1\}$, $I_b = \{2\}$, and $I_c = \emptyset$ for $c \neq a, b$. To simplify writing, we will consider only the case $a > b$. The proof in the case $a < b$ is completely similar.

Let $a > b$. Then $\tilde{U}_I^{(k)} = \tilde{U}_{s(I)}^{(k)} = 1$ for $k < b$, and

$$\tilde{U}_I^{(k)} = \tilde{U}_{s(I)}^{(k)} = \frac{\vartheta(h \mu_{k+1} t_1^{(k+1)} / (\mu_b t_1^{(k)}))}{\vartheta(h \mu_{k+1} / \mu_b)}, \quad b \leq k < a - 1,$$

$$\tilde{U}_I^{(a-1)} = \frac{\vartheta(h t_1^{(a)} / t_1^{(a-1)}) \vartheta(\mu_a t_2^{(a)} / (\mu_b t_1^{(a-1)}))}{\vartheta(\mu_a / \mu_b)},$$

$$\tilde{U}_{s(I)}^{(a-1)} = \frac{\vartheta(t_2^{(a)} / t_1^{(a-1)}) \vartheta(h \mu_a t_1^{(a)} / (\mu_b t_1^{(a-1)}))}{\vartheta(h \mu_a / \mu_b)},$$

$$(2.25) \quad \tilde{U}_I^{(k)} = \zeta(t_1^{(k)}, t_2^{(k)}, t_1^{(k+1)}, t_2^{(k+1)}, \mu_b / \mu_a, \mu_{k+1} / \mu_b), \quad k \geq a,$$

$$(2.26) \quad \tilde{U}_{s(I)}^{(k)} = \zeta(t_1^{(k)}, t_2^{(k)}, t_1^{(k+1)}, t_2^{(k+1)}, \mu_a / \mu_b, \mu_{k+1} / \mu_a), \quad k \geq a,$$

where ζ is defined by (2.24). Set

$$(2.27) \quad \rho(x) = \frac{\vartheta(xh)}{\vartheta(x)}, \quad \varphi(x) = \frac{\vartheta(x \mu_b / \mu_a) \vartheta(h)}{\vartheta(x) \vartheta(\mu_b / \mu_a)}.$$

We will prove the equality

$$(2.28) \quad \begin{aligned} W_I^{(k)}(\mathbf{t}^{(k-1)}, t_1^{(k)}, t_2^{(k)}) &= \\ &= \rho(t_1^{(k)} / t_2^{(k)}) W_{s(I)}^{(k)}(\mathbf{t}^{(k-1)}, t_2^{(k)}, t_1^{(k)}) - \varphi(t_1^{(k)} / t_2^{(k)}) W_{s(I)}^{(k)}(\mathbf{t}^{(k-1)}, t_1^{(k)}, t_2^{(k)}) \end{aligned}$$

for $k \geq a$. By formulae (2.8), (2.14), (2.22), (2.27), this equality proves relation (2.20).

The proof of formula (2.28) is by induction on k . The base of induction is $k = a$, when formula (2.28) reduces to identity (2.6). For the induction step, we multiply formula (2.28)

by $\tilde{U}_I^{(k)}$, see (2.25), and symmetrize over $t_1^{(k)}, t_2^{(k)}$:

$$\begin{aligned} & \text{Sym}_{t_1^{(k)}, t_2^{(k)}} W_I^{(k)}(\mathbf{t}^{(k-1)}, t_1^{(k)}, t_2^{(k)}) \zeta(t_1^{(k)}, t_2^{(k)}, t_1^{(k+1)}, t_2^{(k+1)}, \mu_b/\mu_a, \mu_{k+1}/\mu_b) = \\ & = \text{Sym}_{t_1^{(k)}, t_2^{(k)}} \rho(t_1^{(k)}/t_2^{(k)}) W_{s(I)}^{(k)}(\mathbf{t}^{(k-1)}, t_2^{(k)}, t_1^{(k)}) \zeta(t_1^{(k)}, t_2^{(k)}, t_1^{(k+1)}, t_2^{(k+1)}, \mu_b/\mu_a, \mu_{k+1}/\mu_b) - \\ & \quad - \text{Sym}_{t_1^{(k)}, t_2^{(k)}} \varphi(t_1^{(k)}/t_2^{(k)}) W_{s(I)}^{(k)}(\mathbf{t}^{(k-1)}, t_1^{(k)}, t_2^{(k)}) \zeta(t_1^{(k)}, t_2^{(k)}, t_1^{(k+1)}, t_2^{(k+1)}, \mu_b/\mu_a, \mu_{k+1}/\mu_b). \end{aligned}$$

Now we evaluate the left-hand side by formula (2.23) and swap $t_1^{(k)}, t_2^{(k)}$ in the first summand in the right-hand side to factor out the common factor $W_{s(I)}^{(k)}(\mathbf{t}^{(k-1)}, t_1^{(k)}, t_2^{(k)})$:

$$\begin{aligned} (2.29) \quad & W_I^{(k+1)}(\mathbf{t}^{(k)}, t_1^{(k+1)}, t_2^{(k+1)}) = \\ & = \text{Sym}_{t_1^{(k)}, t_2^{(k)}} \left(W_{s(I)}^{(k)}(\mathbf{t}^{(k-1)}, t_1^{(k)}, t_2^{(k)}) \left(\rho(t_2^{(k)}/t_1^{(k)}) \zeta(t_2^{(k)}, t_1^{(k)}, t_1^{(k+1)}, t_2^{(k+1)}, \mu_b/\mu_a, \mu_{k+1}/\mu_b) - \right. \right. \\ & \quad \left. \left. - \varphi(t_1^{(k)}/t_2^{(k)}) \zeta(t_1^{(k)}, t_2^{(k)}, t_1^{(k+1)}, t_2^{(k+1)}, \mu_b/\mu_a, \mu_{k+1}/\mu_b) \right) \right). \end{aligned}$$

The next step is to use the identity

$$\begin{aligned} & \rho(x_1/x_2) \zeta(x_1, x_2, y_1, y_2, \alpha_1, \alpha_2) - \varphi(x_2/x_1) \zeta(x_2, x_1, y_1, y_2, \alpha_1, \alpha_2) = \\ & = \rho(y_1/y_2) \zeta(x_2, x_1, y_2, y_1, 1/\alpha_1, \alpha_1\alpha_2) - \varphi(y_1/y_2) \zeta(x_2, x_1, y_1, y_2, 1/\alpha_1, \alpha_1\alpha_2), \end{aligned}$$

equivalent to (2.7), to get

$$\begin{aligned} & W_I^{(k+1)}(\mathbf{t}^{(k)}, t_1^{(k+1)}, t_2^{(k+1)}) = \\ & = \text{Sym}_{t_1^{(k)}, t_2^{(k)}} W_{s(I)}^{(k)}(\mathbf{t}^{(k-1)}, t_1^{(k)}, t_2^{(k)}) \rho(t_1^{(k+1)}/t_2^{(k+1)}) \zeta(t_1^{(k)}, t_2^{(k)}, t_2^{(k+1)}, t_1^{(k+1)}, \mu_a/\mu_b, \mu_{k+1}/\mu_a) - \\ & \quad - \text{Sym}_{t_1^{(k)}, t_2^{(k)}} W_{s(I)}^{(k)}(\mathbf{t}^{(k-1)}, t_1^{(k)}, t_2^{(k)}) \varphi(t_1^{(k+1)}/t_2^{(k+1)}) \zeta(t_1^{(k)}, t_2^{(k)}, t_1^{(k+1)}, t_2^{(k+1)}, \mu_a/\mu_b, \mu_{k+1}/\mu_a). \end{aligned}$$

Applying formulae (2.26) and (2.23) in the right-hand side of the last equality transforms it into formula (2.28) with k replaced by $k+1$, which completes the induction step.

Theorem 2.2 is proved. \square

2.6. Transformation properties.

Lemma 2.3. *For $I \in \mathcal{I}_\lambda$, denote*

$$(2.30) \quad G_\lambda(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu}) = \prod_{k=1}^{N-1} \prod_{a=1}^{\lambda^{(k)}} \prod_{c=1}^{\lambda^{(k+1)}} \vartheta(t_c^{(k+1)}/t_a^{(k)}) \prod_{k=1}^{N-1} \frac{\vartheta(t_1^{(k)} \dots t_{\lambda^{(k)}}^{(k)} h^{-\lambda_k} \mu_k/\mu_{k+1})}{\vartheta(t_1^{(k)} \dots t_{\lambda^{(k)}}^{(k)}) \vartheta(h^{-\lambda_k} \mu_k/\mu_{k+1})},$$

$$(2.31) \quad G_I(\mathbf{z}, h, \boldsymbol{\mu}) = (\vartheta(h))^{d_I} \prod_{j=1}^{N-1} \frac{\vartheta(h^{-\lambda_j - \dots - \lambda_{N-1}} \mu_j / \mu_N) \vartheta(\prod_{a \in I_j} z_a)}{\vartheta(h^{-\lambda_j - \dots - \lambda_{N-1}} (\mu_j / \mu_N) \prod_{a \in I_j} z_a)} \\ \times \prod_{j=1}^N \prod_{a \in I_j} \frac{(\vartheta(h))^{(a-1-p_{I,j}(a)-\lambda_1-\dots-\lambda_{j-1})^2} \vartheta(z_a)}{\vartheta(z_a h^{a-1-p_{I,j}(a)-\lambda_1-\dots-\lambda_{j-1}})},$$

where $t_c^{(N)} = z_c$, and $p_{I,j}(a) = |I_j \cap \{1, \dots, a-1\}|$, see (2.12), and

$$d_I = \sum_{j=1}^{N-1} \left(\frac{\lambda^{(j)}(\lambda^{(j)}+1)}{2} + \sum_{a \in I_j} \left((N-j)(a-1) - \sum_{k=j+2}^N (k-j-1)p_{I,k}(a) \right) \right).$$

Set

$$(2.32) \quad G_{\lambda,I}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu}) = G_{\lambda}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu}) G_I(\mathbf{z}, h, \boldsymbol{\mu}),$$

Then the ratio $W_I^{\text{ell}}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu})/G_{\lambda,I}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu})$ does not change if any of the variables $t_i^{(k)}$, h , μ_j is multiplied by q .

Proof. By formula (2.10), the ratio $U_I^{\text{ell}}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu})/G_{\lambda,I}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu})$ does not change if any of the variables $t_i^{(k)}$, h , μ_j is multiplied by q . Since $G_{\lambda,I}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu})$ is symmetric in $t_1^{(i)}, \dots, t_{\lambda^{(i)}}^{(i)}$ for each $i = 1, \dots, N-1$, the same is true for $W_I^{\text{ell}}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu})/G_{\lambda,I}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu})$. \square

Notice that

$$a-1-p_{I,j}(a)-\lambda_1-\dots-\lambda_{j-1} = \\ = |\{1, \dots, a-1\} \cap (I_{j+1} \cup \dots \cup I_N)| - |\{a+1, \dots, n\} \cap (I_1 \cup \dots \cup I_{j-1})|$$

Let $\nu_k = h^{\lambda_1 + \dots + \lambda_{k-1}} \mu_k$. Then $h^{-\lambda_k} \mu_k / \mu_{k+1} = \nu_k / \nu_{k+1}$ and $h^{-\lambda_j - \dots - \lambda_{N-1}} \mu_j / \mu_N = \nu_j / \nu_N$.

Example 1. $N = 2$, $I_1 = \{s\}$, $\lambda_1 = 1$, $\lambda_2 = n-1$.

$$W_I(t, z_1, \dots, z_n) = \vartheta(h) \prod_{a=1}^{s-1} \vartheta(h z_a / t) \frac{\vartheta(h^{2-s} (\mu_2 / \mu_1) z_s / t)}{\vartheta(h^{2-s} \mu_2 / \mu_1)} \prod_{b=s+1}^n \vartheta(z_b / t).$$

The product (2.32) equals

$$(\vartheta(h))^{s^2} \prod_{a=1}^n \vartheta(z_a / t) \frac{\vartheta(h^{-1} t \mu_1 / \mu_2)}{\vartheta(h^{-1} \mu_1 / \mu_2) \vartheta(t)} \frac{\vartheta(h^{-1} \mu_1 / \mu_2) \vartheta(z_s)}{\vartheta(h^{-1} z_s \mu_1 / \mu_2)} \frac{\vartheta(z_s)}{\vartheta(z_s h^{s-1})} \prod_{b=1}^{s-1} \frac{\vartheta(z_b)}{\vartheta(z_b h^{-1})} \prod_{b=s+1}^n \frac{\vartheta(z_b)}{\vartheta(z_b)},$$

and $\nu_1 = \mu_1$, $\nu_2 = h \mu_2$, $h^{-1} \mu_1 / \mu_2 = \nu_1 / \nu_2$.

Example 2. $N = 2$, $n = 3$, $I_1 = \{1, 2\}$, $I_2 = \{3\}$, $\lambda_1 = 2$, $\lambda_2 = 1$.

$$W_I(t_1, t_2, z_1, z_2, z_3) = (\vartheta(h))^2 \vartheta(z_2 / t_1) \vartheta(z_3 / t_1) \vartheta(h z_1 / t_2) \vartheta(z_3 / t_2) \times \\ \times \frac{\vartheta(h (\mu_2 / \mu_1) z_1 / t_1) \vartheta(h^2 (\mu_2 / \mu_1) z_2 / t_2)}{\vartheta(h \mu_2 / \mu_1) \vartheta(h^2 \mu_2 / \mu_1)} \frac{\vartheta(h t_2 / t_1)}{\vartheta(t_2 / t_1)} + [t_1 \leftrightarrow t_2].$$

The product (2.32) equals

$$(\vartheta(h))^4 \prod_{a=1}^2 \prod_{c=1}^3 \vartheta(z_c/t_a) \times \frac{\vartheta(t_1 t_2 h^{-2} \mu_1/\mu_2)}{\vartheta(t_1 t_2) \vartheta(h^{-2} \mu_1/\mu_2)} \times \frac{\vartheta(h^{-2} \mu_1/\mu_2) \vartheta(z_1 z_2)}{\vartheta(h^{-2} z_1 z_2 \mu_1/\mu_2)} \times \frac{\vartheta(z_1)}{\vartheta(z_1)} \frac{\vartheta(z_2)}{\vartheta(z_2)} \frac{\vartheta(z_3)}{\vartheta(z_3)},$$

and $\nu_1 = \mu_1$, $\nu_2 = h^2 \mu_2$, $h^{-2} \mu_1/\mu_2 = \nu_1/\nu_2$.

Trivial fractions from (2.32) are not removed in the Examples.

2.7. Substitution properties of elliptic weight functions. Recall that for $I \in \mathcal{I}_\lambda$ we use the notation $\bigcup_{a=1}^k I_a = \{i_1^{(k)} < \dots < i_{\lambda^{(k)}}^{(k)}\}$. For a function $f(\mathbf{t}, \mathbf{z}, h)$, we denote by $f(\mathbf{z}_I, \mathbf{z}, h)$ the result of the substitution $t_a^{(k)} = z_{i_a^{(k)}}$ for all $k = 1, \dots, N$, $a = 1, \dots, \lambda^{(k)}$.

For any $\sigma \in S_n$, we define the *combinatorial* partial ordering \leq_σ on \mathcal{I}_λ . For $I, J \in \mathcal{I}_\lambda$, let

$$\sigma^{-1}\left(\bigcup_{m=1}^k I_m\right) = \{a_1^{(k)} < \dots < a_{\lambda^{(k)}}^{(k)}\}, \quad \sigma^{-1}\left(\bigcup_{\ell=1}^k J_\ell\right) = \{b_1^{(k)} < \dots < b_{\lambda^{(k)}}^{(k)}\},$$

for $k = 1, \dots, N-1$. We say that $J \leq_\sigma I$ if $b_i^{(k)} \leq a_i^{(k)}$ for all $k = 1, \dots, N-1$, $i = 1, \dots, \lambda^{(k)}$. Notice that $J \leq_\sigma I$ if and only if $\sigma^{-1}(J) \leq_{\text{id}} \sigma^{-1}(I)$.

Set

$$(2.33) \quad E_\lambda^{\text{ell}}(\mathbf{t}, h) = \prod_{k=1}^{N-1} \prod_{a=1}^{\lambda^{(k)}} \prod_{b=1}^{\lambda^{(k)}} \vartheta(ht_b^{(k)}/t_a^{(k)})$$

and

$$(2.34) \quad \tilde{W}_{\sigma, I}^{\text{ell}}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu}) = \frac{W_{\sigma, I}^{\text{ell}}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu})}{E_\lambda^{\text{ell}}(\mathbf{t}, h)}.$$

The proofs of the following three lemmas are straightforward modifications of the proofs in [RTV2, Sect. 6.1]. We indicate main steps of the proof after formulating the lemmas.

Lemma 2.4. *We have $\tilde{W}_{\sigma, I}^{\text{ell}}(\mathbf{z}_J, \mathbf{z}, h, \boldsymbol{\mu}) = 0$ unless $J \leq_\sigma I$.*

Define

$$(2.35) \quad P_{\sigma, I}^{\text{ell}}(\mathbf{z}, h) = \prod_{k < l} \prod_{\sigma(a) \in I_k} \left(\prod_{\substack{\sigma(b) \in I_l \\ b < a}} \vartheta(hz_{\sigma(b)}/z_{\sigma(a)}) \prod_{\substack{\sigma(b) \in I_l \\ b > a}} \vartheta(z_{\sigma(b)}/z_{\sigma(a)}) \right).$$

Lemma 2.5. *We have $\tilde{W}_{\sigma, I}^{\text{ell}}(\mathbf{z}_I, \mathbf{z}, h, \boldsymbol{\mu}) = P_{\sigma, I}^{\text{ell}}(\mathbf{z}, h)$.*

Recall the function $\psi_I(h, \boldsymbol{\mu})$, see (2.13). For $J \in \mathcal{I}_\lambda$, denote

$$(2.36) \quad e_{\sigma, J, -}^{\text{ell, vert}}(\mathbf{z}, h) = \prod_{k < l} \prod_{\sigma(a) \in J_k} \prod_{\substack{\sigma(b) \in J_l \\ b < a}} \vartheta(hz_{\sigma(b)}/z_{\sigma(a)}).$$

Lemma 2.6. *For all $I, J \in \mathcal{I}_\lambda$, the function $\psi_I(h, \boldsymbol{\mu}) \tilde{W}_{\sigma, I}^{\text{ell}}(\mathbf{z}_J, \mathbf{z}, h, \boldsymbol{\mu}) / e_{\sigma, J, -}^{\text{ell, vert}}(\mathbf{z}, h)$ lifts to a regular function on the universal cover of $(\mathbb{C}^\times)^{n+1+N}$.*

The universal cover in the last lemma is due to presence of the square root $x^{1/2}$ in formula (2.1) for $\vartheta(x)$.

Proof of Lemmas 2.4–2.6. By formula (2.14), it suffices to verify the claims for σ being the identity permutation. The function $W_{\text{id},I}^{\text{ell}}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu})$ is the symmetrization of product (2.10). By inspection of formula (2.10), each term in the symmetrization sum acquire a zero factor at $\mathbf{t} = \mathbf{z}_J$ unless $J \leq_{\text{id}} I$, and $E_{\lambda}^{\text{ell}}(\mathbf{z}_J, h) \neq 0$. For $J = I$, the only term of the sum for $W_{\text{id},I}^{\text{ell}}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu})$ that does not vanish at $\mathbf{t} = \mathbf{z}_J$ corresponds to the identity element, and equals $U_I^{\text{ell}}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu})$. Then $\tilde{W}_{\text{id},I}^{\text{ell}}(\mathbf{z}_I, \mathbf{z}, h, \boldsymbol{\mu}) = U_{\text{id},I}^{\text{ell}}(\mathbf{z}_I, \mathbf{z}, h, \boldsymbol{\mu})/E_{\lambda}^{\text{ell}}(\mathbf{z}_I, h) = P_{\text{id},I}^{\text{ell}}(\mathbf{z}, h)$. Finally for $J <_{\text{id}} I$, inspection of (2.10) shows that each potentially nonzero term of the sum for $W_{\text{id},I}^{\text{ell}}(\mathbf{z}_J, \mathbf{z}, h, \boldsymbol{\mu})$ is divisible by $E_{\lambda}^{\text{ell}}(\mathbf{z}_I, h)e_{\sigma, J, -}^{\text{ell,vert}}(\mathbf{z}, h)$. \square

2.8. Orthogonality. Let $\sigma_0 \in S_n$ be the longest permutation in S_n . Denote $h^\lambda \boldsymbol{\mu}^{-1} = (h^{\lambda_1} \mu_1^{-1}, \dots, h^{\lambda_N} \mu_N^{-1})^{-1}$. Let $\tilde{\mathbf{t}}$ be an additional set of variables similar to \mathbf{t} . Define a function $\Xi(\mathbf{t}, \tilde{\mathbf{t}}, \mathbf{z}, h, \boldsymbol{\mu})$,

$$(2.37) \quad \Xi(\mathbf{t}, \tilde{\mathbf{t}}, \mathbf{z}, h, \boldsymbol{\mu}) = \sum_{I \in \mathcal{I}_\lambda} \tilde{W}_{\text{id},I}^{\text{ell}}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu}) \tilde{W}_{\sigma_0, I}^{\text{ell}}(\tilde{\mathbf{t}}, \mathbf{z}, h, h^\lambda \boldsymbol{\mu}^{-1}).$$

Proposition 2.7. *The function $\Xi(\mathbf{t}, \tilde{\mathbf{t}}, \mathbf{z}, h, \boldsymbol{\mu})$ is symmetric in \mathbf{z} .*

Proof. The statement follows from Theorem 2.2. \square

Proposition 2.8. *For $I, J \in \mathcal{I}_\lambda$, we have*

$$(2.38) \quad \Xi(\mathbf{z}_I, \mathbf{z}_J, \mathbf{z}, h, \boldsymbol{\mu}) = R^{\text{ell}}(\mathbf{z}_I) Q^{\text{ell}}(\mathbf{z}_I, h) \delta_{I,J},$$

where

$$(2.39) \quad R^{\text{ell}}(\mathbf{z}_I) = \prod_{k < l} \prod_{a \in I_k} \prod_{b \in I_l} \vartheta(z_b/z_a), \quad Q^{\text{ell}}(\mathbf{z}_I, h) = \prod_{k < l} \prod_{a \in I_k} \prod_{b \in I_l} \vartheta(hz_b/z_a).$$

Proof. Denote by $f_{I,J}(\mathbf{z})$ one of the sides of formula (2.38). Then either by Proposition 2.7 or by formulae (2.39), we have $f_{\sigma(I), \sigma(J)}(\mathbf{z}) = f_{I,J}(z_{\sigma(1)}, \dots, z_{\sigma(n)})$ for any $\sigma \in S_n$. Thus it is enough to verify formula (2.38) for $J = (\{1, \dots, \lambda_1\}, \dots, \{n - \lambda_n + 1, \dots, n\})$. In this case, $I \leq_{\sigma_0} J$ for any $I \in \mathcal{I}_\lambda$, and formula (2.38) follows from Lemmas 2.4, 2.5. \square

Theorem 2.9. *For $J, K \in \mathcal{I}_\lambda$, we have*

$$(2.40) \quad \sum_{I \in \mathcal{I}_\lambda} \frac{\tilde{W}_{\text{id},J}^{\text{ell}}(\mathbf{z}_I, \mathbf{z}, h, \boldsymbol{\mu}) \tilde{W}_{\sigma_0, K}^{\text{ell}}(\mathbf{z}_I, \mathbf{z}, h, h^\lambda \boldsymbol{\mu}^{-1})}{R^{\text{ell}}(\mathbf{z}_I) Q^{\text{ell}}(\mathbf{z}_I, h)} = \delta_{J,K}.$$

Proof. Consider the matrices $\widehat{W}, \widehat{W}', \widehat{Q}$ with entries

$$\widehat{W}_{I,J} = \tilde{W}_{\text{id},J}^{\text{ell}}(\mathbf{z}_I, \mathbf{z}, h, \boldsymbol{\mu}), \quad \widehat{W}'_{K,I} = \tilde{W}_{\sigma_0, K}^{\text{ell}}(\mathbf{z}_I, \mathbf{z}, h, h^\lambda \boldsymbol{\mu}^{-1}), \quad \widehat{Q}_{I,J} = R^{\text{ell}}(\mathbf{z}_I) Q^{\text{ell}}(\mathbf{z}_I, h) \delta_{I,J}.$$

Formula (2.38) yields a matrix equality $\widehat{W} \widehat{W}' = \widehat{Q}$. Therefore, $\widehat{W}' \widehat{Q}^{-1} \widehat{W} = 1$, which is exactly formula (2.40). \square

Remark. Denote $\mathbf{t}^{-1} = ((t_1^{(1)})^{-1}, \dots, (t_{\lambda^{(N-1)}}^{(N-1)})^{-1})$, $\mathbf{z}^{-1} = (z_1^{-1}, \dots, z_n^{-1})$. Notice that all the consideration is essentially invariant under the simultaneous replacement

$$h \rightarrow h^{-1}, \quad \mathbf{z} \rightarrow \mathbf{z}^{-1}, \quad \mathbf{t} \rightarrow \mathbf{t}^{-1}, \quad \boldsymbol{\mu} \rightarrow \boldsymbol{\mu}^{-1},$$

for instance, $\mathcal{R}(x, h, \boldsymbol{\mu}) = \mathcal{R}(x^{-1}, h^{-1}, \boldsymbol{\mu}^{-1})$,

$$W_{\sigma, I}^{\text{ell}}(\mathbf{t}^{-1}, \mathbf{z}^{-1}, h^{-1}, \boldsymbol{\mu}^{-1}) = (-1)^{\sum_{k=1}^{N-1} \lambda^{(k)} \lambda^{(k+1)}} W_{\sigma, I}^{\text{ell}}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu}).$$

3. TRIGONOMETRIC WEIGHT FUNCTIONS

3.1. Alcoves. Consider \mathbb{R}^N with coordinates $\nu = (\nu_1, \dots, \nu_N)$. A *wall* in \mathbb{R}^N is a hyperplane defined by an equation of the form $\nu_j - \nu_i = m$, $1 \leq i < j \leq N$, $m \in \mathbb{Z}$. Connected components of the complement in \mathbb{R}^N to the union of all walls are called *alcoves*. Let Δ be an alcove and $\nu \in \Delta$. For $1 \leq i < j \leq N$, denote $m_{i,j} = \lfloor \nu_j - \nu_i \rfloor \in \mathbb{Z}$, that is, $0 < \nu_j - \nu_i - m_{i,j} < 1$. The numbers $(m_{i,j})_{1 \leq i < j \leq N}$ depend on the alcove Δ , but not on $\nu \in \Delta$; they are called the *integer characteristics* of the alcove Δ . The *anti-dominant* alcove is the alcove with zero integer characteristics.

Let Δ be an alcove with integer characteristics $(m_{i,j})$. Recall the $\mathbf{t}, \mathbf{z}, h$ variables from Section 2.4. For $I \in \mathcal{I}_\lambda$, define the *trigonometric weight function*

$$(3.1) \quad W_I^\Delta(\mathbf{t}, \mathbf{z}, h) = (1-h)^{\lambda^{\{1\}}} \text{Sym}_{t^{(1)}} \dots \text{Sym}_{t^{(N-1)}} U_I^\Delta,$$

where

$$\text{Sym}_{t^{(k)}} f(t_1^{(k)}, \dots, t_{\lambda^{(k)}}^{(k)}) = \sum_{\sigma \in S_{\lambda^{(k)}}} f(t_{\sigma(1)}^{(k)}, \dots, t_{\sigma(\lambda^{(k)})}^{(k)}),$$

$$(3.2) \quad U_I^\Delta = \prod_{k=1}^{N-1} \prod_{a=1}^{\lambda^{(k)}} \left(\prod_{c=1}^{\lambda^{(k+1)}} \psi_{I, k, a, c}^\Delta (t_c^{(k+1)} / t_a^{(k)}) \prod_{b=a+1}^{\lambda^{(k)}} \frac{1 - ht_b^{(k)} / t_a^{(k)}}{1 - t_b^{(k)} / t_a^{(k)}} \right),$$

$$(3.3) \quad \psi_{I, k, a, c}^\Delta(x) = \begin{cases} 1 - hx, & \text{if } i_c^{(k+1)} < i_a^{(k)}, \\ x^{-m_{j(I, k, a), k+1}}, & \text{if } i_c^{(k+1)} = i_a^{(k)}, \\ 1 - x, & \text{if } i_c^{(k+1)} > i_a^{(k)}. \end{cases}$$

where $j(I, k, a) \in \{1, \dots, N\}$ is such that $i_a^{(k)} \in I_{j(I, k, a)}$, cf. (2.11).

For $\sigma \in S_n$ and $I \in \mathcal{I}_\lambda$, define the trigonometric weight function

$$(3.4) \quad W_{\sigma, I}^\Delta(\mathbf{t}, \mathbf{z}, h) = W_{\sigma^{-1}(I)}^\Delta(\mathbf{t}, z_{\sigma(1)}, \dots, z_{\sigma(n)}, h),$$

where $\sigma^{-1}(I) = (\sigma^{-1}(I_1), \dots, \sigma^{-1}(I_N))$. Hence, $W_I^\Delta = W_{\text{id}, I}^\Delta$.

Define

$$(3.5) \quad E(\mathbf{t}, h) = \prod_{k=1}^{N-1} \prod_{a=1}^{\lambda^{(k)}} \prod_{b=1}^{\lambda^{(k)}} \left(1 - \frac{ht_b^{(k)}}{t_a^{(k)}} \right)$$

and

$$(3.6) \quad \tilde{W}_{\sigma, I}^\Delta(\mathbf{t}, \mathbf{z}, h) = \frac{W_{\sigma, I}^\Delta(\mathbf{t}, \mathbf{z}, h)}{E(\mathbf{t}, h)}.$$

Remark. The trigonometric weight functions of [RTV2] correspond to $W_{\sigma, I}^{\Delta}(\mathbf{t}, \mathbf{z}, h)$ with the anti-dominant alcove Δ .

3.2. Elliptic versus trigonometric weight functions: the $q \rightarrow 0$ limit. Let Δ be an alcove with integer characteristics $(m_{j,k})$, and let $\nu \in \Delta$. Define

$$(3.7) \quad \mu_j = e^{2\pi i \tau \nu_j}, \quad j = 1, \dots, N.$$

Then for $1 \leq j < k \leq N$, we have $\mu_k/\mu_j = q^{m_{j,k} + \varepsilon_{j,k}}$, where $\varepsilon_{j,k} = \nu_k - \nu_j - m_{j,k} \in (0, 1)$.

For $I \in \mathcal{I}_{\lambda}$, define

$$(3.8) \quad \ell_I = \#\{(a, b) \in \{1, \dots, n\}^2 \mid a \in I_j, b \in I_k, a > b, j < k\}.$$

Equivalently, ℓ_I is the length of the permutation $\sigma_I \in S_n$ such that

$$(3.9) \quad I_j = \{\sigma_I(\lambda^{(j-1)} + 1) < \dots < \sigma_I(\lambda^{(j)})\}, \quad j = 1, \dots, N.$$

Theorem 3.1. *Let $h, \mathbf{z}, \mathbf{t}$ be fixed. As $q \rightarrow 0$, we have*

$$(3.10) \quad W_I^{\text{ell}}(\mathbf{t}, \mathbf{z}, h, \boldsymbol{\mu}) \rightarrow c(h) h^{-\ell_I/2} (X(\mathbf{t}))^{-1/2} W_I^{\Delta}(\mathbf{t}, \mathbf{z}, h),$$

where $c(h) = (-1)^{\sum_{k=1}^{N-1} \lambda^{(k)} \lambda^{(k+1)}} h^{-\sum_{k=1}^{N-1} (\lambda^{(k)})^2/2}$ and

$$X(\mathbf{t}) = \prod_{k=1}^{N-1} \prod_{a=1}^{\lambda^{(k)}} \prod_{b=1}^{\lambda^{(k+1)}} (t_b^{(k+1)}/t_a^{(k)}) = \prod_{k=1}^{N-1} \prod_{a=1}^{\lambda^{(k)}} (t_a^{(k)})^{-\lambda_k - \lambda_{k+1}} \prod_{a=1}^n z_a^{n - \lambda_N}$$

Proof. The statement follows from formulae (2.9)–(2.12) and (3.1)–(3.3) for the weight functions, and the limit properties (2.5). The next formulae simplify the expression for $c(h)$:

$$\sum_{a=1}^{\lambda^{(k)}} \sum_{c=1}^k p_{I,c}(i_a^{(k)}) = \frac{\lambda^{(k)} (\lambda^{(k)} - 1)}{2}, \quad \sum_{k=1}^{N-1} \sum_{a=1}^{\lambda^{(k)}} p_{I,k+1}(i_a^{(k)}) = \ell_I. \quad \square$$

Remark. In the geometric language of Section 4, ℓ_I equals the dimension of the Schubert cell $\Omega_{\text{id}, I}$, see (4.1), (4.2). Thus the ratio of the factor $h^{-\ell_I/2}$ in (3.10) and the factor $h^{\text{codim}(\Omega_{\text{id}, I} \subset \mathcal{F}_{\lambda})/2}$ in the normalization condition in Section 5 equals $h^{-(\dim \mathcal{F}_{\lambda})/2}$ and does not depend on $I \in \mathcal{I}_{\lambda}$.

Remark. Notice the substitution formula, cf. (3.15):

$$X(\mathbf{z}_I) = \prod_{1 \leq j < k \leq N} \prod_{a \in I_j} \prod_{b \in I_k} (z_b/z_a).$$

Remark. Theorem 3.1 is our attempt to interpret [AO, Proposition 3.5] as a property of the elliptic and trigonometric weight functions. In [AO, Proposition 3.5] the authors claim for a Nakajima variety that an elliptic stable envelope converges to a certain K -theoretic stable envelope under a certain limit. The K -theoretic stable envelope are not precisely defined in [AO] and the proof of Proposition 3.5 in [AO] is only sketched.

3.3. Recursion properties. Let $s_{a,a+1}(\mathbf{z}) = (z_1, \dots, z_{a-1}, z_{a+1}, z_a, z_{a+2}, \dots, z_n)$.

Theorem 3.2. *Consider an alcove Δ with integer characteristics $(m_{i,j})$, and $I \in \mathcal{I}_\lambda$ such that $a \in I_k$, $a+1 \in I_l$. If $k = l$, then*

$$(3.11) \quad W_I^\Delta(\mathbf{t}, s_{a,a+1}(\mathbf{z}), h) = W_I^\Delta(\mathbf{t}, \mathbf{z}, h).$$

If $k < l$, then

$$(3.12) \quad \begin{aligned} W_{s_{a,a+1}(I)}^\Delta(\mathbf{t}, \mathbf{z}, h) &= \\ &= \frac{1 - h z_a / z_{a+1}}{1 - z_a / z_{a+1}} W_I^\Delta(\mathbf{t}, s_{a,a+1}(\mathbf{z}), h) + (h - 1) \frac{(z_a / z_{a+1})^{m_{k,l}+1}}{1 - z_a / z_{a+1}} W_I^\Delta(\mathbf{t}, \mathbf{z}, h). \end{aligned}$$

If $k > l$, then

$$(3.13) \quad \begin{aligned} W_{s_{a,a+1}(I)}^\Delta(\mathbf{t}, \mathbf{z}, h) &= \\ &= \frac{1 - h^{-1} z_{a+1} / z_a}{1 - z_{a+1} / z_a} W_I^\Delta(\mathbf{t}, s_{a,a+1}(\mathbf{z}), h) + (h^{-1} - 1) \frac{(z_{a+1} / z_a)^{m_{l,k}+1}}{1 - z_{a+1} / z_a} W_I^\Delta(\mathbf{t}, \mathbf{z}, h). \end{aligned}$$

Proof. The statement follows from Theorems 2.2 and 3.1. \square

3.4. Substitution properties of trigonometric weight functions. Recall that for $I \in \mathcal{I}_\lambda$ we use the notation $\bigcup_{a=1}^k I_a = \{i_1^{(k)} < \dots < i_{\lambda^{(k)}}^{(k)}\}$. For a function $f(\mathbf{t}, \mathbf{z}, h)$, we denote by $f(\mathbf{z}_I, \mathbf{z}, h)$ the result of the substitution $t_a^{(k)} = z_{i_a^{(k)}}$ for all $k = 1, \dots, N$, $a = 1, \dots, \lambda^{(k)}$. Recall the partial ordering \leq_σ on \mathcal{I}_λ defined in Section 2.7.

The proofs of the following four lemmas are straightforward modifications of those in [RTV2, Section 6.1], cf. the proof of Lemmas 2.4–2.6 in Section 2.7.

Lemma 3.3. *For any $\sigma \in S_n$ and $I, J \in \mathcal{I}_\lambda$, the function $\tilde{W}_{\sigma,I}^\Delta(\mathbf{z}_J, \mathbf{z}, h)$ is a Laurent polynomial.*

Lemma 3.4. *We have $\tilde{W}_{\sigma,I}^\Delta(\mathbf{z}_J, \mathbf{z}, h) = 0$ unless $J \leq_\sigma I$.*

Define

$$(3.14) \quad P_{\sigma,I} = \prod_{k < l} \prod_{\sigma(a) \in I_k} \prod_{\substack{\sigma(b) \in I_l \\ b > a}} (-z_{\sigma(b)} / z_{\sigma(a)}).$$

Using definitions (4.7), (4.8) of $e_{\sigma,I} = e_{\sigma,I,-}^{\text{hor}} e_{\sigma,I,-}^{\text{vert}}$, observe that

$$(3.15) \quad P_{\sigma,I} \cdot e_{\sigma,I} = \prod_{k < l} \prod_{\sigma(a) \in I_k} \left(\prod_{\substack{\sigma(b) \in I_l \\ b < a}} (1 - h z_{\sigma(b)} / z_{\sigma(a)}) \prod_{\substack{\sigma(b) \in I_l \\ b > a}} (1 - z_{\sigma(b)} / z_{\sigma(a)}) \right).$$

Lemma 3.5. *We have $\tilde{W}_{\sigma,I}^\Delta(\mathbf{z}_I, \mathbf{z}, h) = P_{\sigma,I} \cdot e_{\sigma,I}$.*

Lemma 3.6. *For any $I, J \in \mathcal{I}_\lambda$, the ratio $\tilde{W}_{\sigma,I}^\Delta(\mathbf{z}_J, \mathbf{z}, h) / e_{\sigma,J,-}^{\text{vert}}$ is a Laurent polynomial.*

3.5. Newton polytope properties. For $a = 1, \dots, n-1$, denote by K_a the automorphism of $\mathbb{C}[z^{\pm 1}, h^{\pm 1}]$ switching z_a and z_{a+1} . That is $K_a(f(\mathbf{z})) = f(s_{a,a+1}(\mathbf{z}))$.

Let $\nu \in \mathbb{R}^N$ belong to the alcove Δ with integer characteristics $(m_{i,j})$. We will use the notation $\varepsilon_{i,j} = \nu_j - \nu_i - m_{i,j}$ for the fractional part of $\nu_j - \nu_i$ for $1 \leq i < j \leq N$; we have $0 < \varepsilon_{i,j} < 1$. Define

$$S_I = S_I(\mathbf{z}) = \prod_{k=1}^N \prod_{a \in I_k} z_a^{-\nu_k} \quad \text{and} \quad \bar{W}_{I,J}^\Delta = \bar{W}_{I,J}^\Delta(\mathbf{z}, h) = W_I^\Delta(\mathbf{z}_J, \mathbf{z}, h) \cdot S_I(\mathbf{z}).$$

Although the notation does not record it, both S_I and $\bar{W}_{I,J}^\Delta$ depend on ν itself, not only on its alcove Δ .

Lemma 3.7. *Let $I \in \mathcal{I}_\lambda$ be such that $a \in I_k, a+1 \in I_l$. If $k = l$, then*

$$(3.16) \quad \bar{W}_{I, s_{a,a+1}(J)}^\Delta = K_a(\bar{W}_{I,J}^\Delta).$$

If $k < l$, then

$$(3.17) \quad \bar{W}_{I, s_{a,a+1}(J)}^\Delta = K_a \left(\frac{1 - z_a/z_{a+1}}{1 - h z_a/z_{a+1}}; \bar{W}_{s_{a,a+1}(I), J}^\Delta + (1-h) \frac{(z_a/z_{a+1})^{1-\varepsilon_{k,l}}}{1 - h z_a/z_{a+1}} \bar{W}_{I,J}^\Delta \right).$$

If $k > l$, then

$$(3.18) \quad \bar{W}_{I, s_{a,a+1}(J)}^\Delta = K_a \left(\frac{1 - z_{a+1}/z_a}{1 - h^{-1} z_{a+1}/z_a} \bar{W}_{s_{a,a+1}(I), J}^\Delta + (1-h^{-1}) \frac{(z_{a+1}/z_a)^{1-\varepsilon_{l,k}}}{1 - h^{-1} z_{a+1}/z_a} \bar{W}_{I,J}^\Delta \right).$$

Proof. Let $k = l$. Then taking into account (3.11),

$$\begin{aligned} K_a(\bar{W}_{I,J}^\Delta) &= K_a(W_I^\Delta(\mathbf{z}_J, \mathbf{z}, h)) K_a(S_I) = W_I^\Delta(\mathbf{z}_{s_{a,a+1}(J)}, s_{a,a+1}(\mathbf{z}), h) S_I = \\ &= W_I^\Delta(\mathbf{z}_{s_{a,a+1}(J)}, \mathbf{z}, h) S_I = \bar{W}_{I, s_{a,a+1}(J)}^\Delta. \end{aligned}$$

Let $k < l$. Formula (3.12), rearranged, gives

$$W_I^\Delta(\mathbf{t}, s_{a,a+1}(\mathbf{z}), h) = \frac{1 - z_a/z_{a+1}}{1 - h z_a/z_{a+1}} W_{s_{a,a+1}(I)}^\Delta(\mathbf{t}, \mathbf{z}, h) + (1-h) \frac{(z_a/z_{a+1})^{m_{k,l}+1}}{1 - h z_a/z_{a+1}} W_I^\Delta(\mathbf{t}, \mathbf{z}, h).$$

Substitute now $\mathbf{t} = \mathbf{z}_J$. Then,

$$\begin{aligned} K_a(W_I^\Delta(\mathbf{z}_{s_{a,a+1}(J)}, \mathbf{z}, h)) &= W_I^\Delta(\mathbf{z}_J, s_{a,a+1}(\mathbf{z}), h) = \\ &= \frac{1 - z_a/z_{a+1}}{1 - h z_a/z_{a+1}} W_{s_{a,a+1}(I)}^\Delta(\mathbf{z}_J, \mathbf{z}, h) + (1-h) \frac{(z_a/z_{a+1})^{m_{k,l}+1}}{1 - h z_a/z_{a+1}} W_I^\Delta(\mathbf{z}_J, \mathbf{z}, h), \end{aligned}$$

that is,

$$\frac{K_a(\bar{W}_{I, s_{a,a+1}(J)}^\Delta)}{K_a(S_I)} = \frac{1 - z_a/z_{a+1}}{1 - h z_a/z_{a+1}} \cdot \frac{\bar{W}_{s_{a,a+1}(I), J}^\Delta}{S_{s_{a,a+1}(I)}} + (1-h) \frac{(z_a/z_{a+1})^{m_{k,l}+1}}{1 - h z_a/z_{a+1}} \cdot \frac{\bar{W}_{I,J}^\Delta}{S_I}.$$

Since $K_a(S_I) = S_{s_{a,a+1}(I)} = (z_{a+1}/z_a)^{\nu_l - \nu_k} S_I = (z_{a+1}/z_a)^{m_{k,l} + \varepsilon_{k,l}} S_I$, we obtain

$$(3.19) \quad K_a(\overline{W}_{I,s_{a,a+1}(J)}^\Delta) = \frac{1 - z_a/z_{a+1}}{1 - h z_a/z_{a+1}} \overline{W}_{s_{a,a+1}(I),J}^\Delta + (1 - h) \frac{(z_a/z_{a+1})^{1 - \varepsilon_{k,l}}}{1 - h z_a/z_{a+1}} \overline{W}_{I,J}^\Delta,$$

which is equivalent to formula (3.17). The case $k > l$ is proved similarly. \square

For a Laurent polynomial $f \in \mathbb{C}[z_1^{\pm 1}, \dots, z_n^{\pm 1}, h^{\pm 1}]$, let $\mathcal{N}(f)$ be the convex hull of the finite set

$$\{(k_1, \dots, k_n) \in \mathbb{Z}^n \mid \text{the coefficient of } z_1^{k_1} \dots z_n^{k_n} \text{ in } f \text{ is not } 0\}.$$

That is, in Newton polytope considerations we treat h as a number, not as a variable. Let $O_{I,J} = \mathcal{N}(\overline{W}_{I,J}^\Delta(z, h))$. The operators K_a act naturally on Newton polytopes, satisfying $K_a(\mathcal{N}(f(z))) = \mathcal{N}(f(s_{a,a+1}(z)))$.

Lemma 3.8. *We have $O_{s_{a,a+1}(J), s_{a,a+1}(J)} = K_a(O_{J,J})$.*

Proof. Let $a \in J_k$, $a + 1 \in J_l$. If $k = l$, then the statement follows from formula (3.16).

Let $k \neq l$. Replacing J by $s_{a,a+1}(J)$ if necessary, we may assume that $k > l$. Denote $I = s_{a,a+1}(J)$. We have $J \not\leq_{\text{id}} I$, hence $\overline{W}_{I,J}^\Delta = 0$, and formula (3.18) yields

$$(1 - h^{-1} z_a/z_{a+1}) \overline{W}_{s_{a,a+1}(J), s_{a,a+1}(J)}^\Delta = (1 - z_a/z_{a+1}) K_a(\overline{W}_{J,J}^\Delta).$$

Taking the Newton polytope of both sides, we obtain

$$T + O_{s_{a,a+1}(J), s_{a,a+1}(J)} = T + K_a(O_{J,J}),$$

where $+$ is the Minkowski sum and $T = \mathcal{N}(1 - h^{-1} z_a/z_{a+1}) = \mathcal{N}(1 - z_a/z_{a+1})$. Using the cancellation law of Minkowski sum, we obtain the statement of the lemma. \square

The following theorem is one of the main results of this paper.

Theorem 3.9. *We have $O_{I,J} \subset O_{J,J}$ for any $I, J \in \mathcal{I}_\lambda$.*

Proof. We will prove by induction on J that $O_{I,J} \subset O_{J,J}$ for all $I \in \mathcal{I}_\lambda$. The induction employs the partial ordering \leq_{id} . The base of induction is

$$J = I^{\max} = (\{n - \lambda_1 + 1, \dots, n\}, \{n - \lambda_1 - \lambda_2 + 1, \dots, n - \lambda_1\}, \dots, \{1, \dots, \lambda_N\}).$$

Then $I \leq_{\text{id}} J$ for all $I \in \mathcal{I}_\lambda$. Hence by Lemma 3.4, $\overline{W}_{I,J}^\Delta = 0$ for $I \neq J$, and the statement holds.

If $J \neq I^{\max}$, let a be such that $J <_{\text{id}} s_{a,a+1}(J) = \tilde{J}$. Let $I \in \mathcal{I}_\lambda$ and define k, l by the rule: $a \in I_k$, $a + 1 \in I_l$. If $k = l$, then

$$O_{I,J} = K_a(O_{I,\tilde{J}}) \subset K_a(O_{\tilde{J},\tilde{J}}) = O_{J,J},$$

where the first equality follows from formula (3.18), the containment holds by the induction assumption, and the last equality is due to Lemma 3.8.

If $k < l$, let O' be the Newton polytope of

$$K_a(\overline{W}_{I,s_{a,a+1}(J)}^\Delta) = \frac{1 - z_a/z_{a+1}}{1 - h z_a/z_{a+1}} \overline{W}_{s_{a,a+1}(I),J}^\Delta + (1 - h) \frac{(z_a/z_{a+1})^{1 - \varepsilon_{k,l}}}{1 - h z_a/z_{a+1}} \overline{W}_{I,J}^\Delta,$$

see (3.19). Let $T = \mathcal{N}(1 - z_a/z_{a+1}) = \mathcal{N}(1 - hz_a/z_{a+1})$ (an interval), and $p = \mathcal{N}((z_a/z_{a+1})^{1-\varepsilon_{k,l}})$ (one point). From the definition of O' we obtain

$$T + O' = \text{c.h.} (T + \mathcal{N}(\overline{W}_{s_{a,a+1}(I), \bar{j}}^\Delta), p + \mathcal{N}(\overline{W}_{I, \bar{j}}^\Delta)),$$

where c.h. stands for convex hull, and $+$ for the Minkowski sum. Since $1 - \varepsilon_{k,l} \in (0, 1)$, we have $p \subset T$, and therefore,

$$T + O' \subset \text{c.h.} (T + \mathcal{N}(\overline{W}_{s_{a,a+1}(I), \bar{j}}^\Delta), T + \mathcal{N}(\overline{W}_{I, \bar{j}}^\Delta)) = T + \text{c.h.} (\mathcal{N}(\overline{W}_{s_{a,a+1}(I), \bar{j}}^\Delta), \mathcal{N}(\overline{W}_{I, \bar{j}}^\Delta)).$$

By the induction assumption, both $\mathcal{N}(\overline{W}_{s_{a,a+1}(I), \bar{j}}^\Delta)$ and $\mathcal{N}(\overline{W}_{I, \bar{j}}^\Delta)$ are contained in $O_{\bar{j}, \bar{j}}$. Thus $T + O' \subset T + O_{\bar{j}, \bar{j}}$, which by cancellation law yields $O' \subset O_{\bar{j}, \bar{j}}$. Therefore,

$$O_{I, J} = K_a(O') \subset K_a(O_{\bar{j}, \bar{j}}) = O_{J, J},$$

where the first equality follows from formula (3.17) and the last equality is due to Lemma 3.8.

The case $k > l$ is proved similarly. \square

Example. For $N = 2$, $\lambda = (1, 2)$, let

$$I_1 = (\{1\}, \{2, 3\}), \quad I_2 = (\{2\}, \{1, 3\}), \quad I_3 = (\{3\}, \{1, 2\}),$$

and let $(\nu_1, \nu_2) \in \mathbb{R}^2$ be such that $\nu_2 - \nu_1 = m_{1,2} + \varepsilon_{1,2}$, where $\varepsilon_{1,2} \in (0, 1)$. Then we have

$$W_{I_1}^\Delta = (1 - h)(t/z_1)^{m_{1,2}}(1 - z_2/t)(1 - z_3/t),$$

$$W_{I_2}^\Delta = (1 - h)(t/z_2)^{m_{1,2}}(1 - hz_1/t)(1 - z_3/t),$$

$$W_{I_3}^\Delta = (1 - h)(t/z_3)^{m_{1,2}}(1 - hz_1/t)(1 - hz_2/t).$$

Hence

$$O_{I_1, I_1} = \mathcal{N}((1 - h)(1 - z_2/z_1)(1 - z_3/z_1) \cdot z_1^{-\nu_1}(z_2 z_3)^{-\nu_2}),$$

$$O_{I_2, I_1} = \mathcal{N}((1 - h)^2(z_1/z_2)^{m_{1,2}}(1 - z_3/z_1) \cdot z_2^{-\nu_1}(z_1 z_3)^{-\nu_2}),$$

$$O_{I_3, I_1} = \mathcal{N}((1 - h)^2(z_1/z_3)^{m_{1,2}}(1 - hz_2/z_1) \cdot z_3^{-\nu_1}(z_1 z_2)^{-\nu_2}).$$

Therefore, the statements $O_{I_2, I_1} \subset O_{I_1, I_1}$ and $O_{I_3, I_1} \subset O_{I_1, I_1}$ after arranging the \mathbf{z} -monomials to the left-hand sides reduce to the statements

$$\mathcal{N}((1 - h)^2(1 - z_3/z_1)(z_2/z_1)^{\varepsilon_{1,2}}) \subset \mathcal{N}((1 - h)(1 - z_2/z_1)(1 - z_3/z_1))$$

and

$$\mathcal{N}((1 - h)^2(1 - hz_2/z_1)(z_3/z_1)^{\varepsilon_{1,2}}) \subset \mathcal{N}((1 - h)(1 - z_2/z_1)(1 - z_3/z_1)),$$

respectively, see Figure 1.




FIGURE 1. $\mathcal{N}((1 - z_3/z_1)(z_2/z_1)^{\epsilon_{1,2}})$, $\mathcal{N}((1 - hz_2/z_1)(z_3/z_1)^{\epsilon_{1,2}})$ are contained in $\mathcal{N}((1 - z_2/z_1)(1 - z_3/z_1))$.

4. PARTIAL FLAG VARIETIES

4.1. **Definitions.** Fix natural numbers N, n . Let $\boldsymbol{\lambda} \in \mathbb{Z}_{\geq 0}^N$, $|\boldsymbol{\lambda}| = \lambda_1 + \dots + \lambda_N = n$. Consider the partial flag variety $\mathcal{F}_{\boldsymbol{\lambda}}$ parametrizing chains of subspaces

$$0 = F_0 \subset F_1 \subset \dots \subset F_N = \mathbb{C}^n$$

with $\dim F_i/F_{i-1} = \lambda_i$, $i = 1, \dots, N$. Denote by $T^*\mathcal{F}_{\boldsymbol{\lambda}}$ the cotangent bundle of $\mathcal{F}_{\boldsymbol{\lambda}}$, and let $\pi : T^*\mathcal{F}_{\boldsymbol{\lambda}} \rightarrow \mathcal{F}_{\boldsymbol{\lambda}}$ be the projection of the bundle.

Let $I = (I_1, \dots, I_N)$ be a partition of $\{1, \dots, n\}$ into disjoint subsets I_1, \dots, I_N . Denote $\mathcal{I}_{\boldsymbol{\lambda}}$ the set of all partitions I with $|I_j| = \lambda_j$, $j = 1, \dots, N$.

Let $\epsilon_1, \dots, \epsilon_n$ be the standard basis of \mathbb{C}^n . For any $I \in \mathcal{I}_{\boldsymbol{\lambda}}$, let $x_I \in \mathcal{F}_{\boldsymbol{\lambda}}$ be the point corresponding to the coordinate flag $F_1 \subset \dots \subset F_N$, where F_i is the span of the standard basis vectors $\epsilon_j \in \mathbb{C}^n$ with $j \in I_1 \cup \dots \cup I_i$. We embed $\mathcal{F}_{\boldsymbol{\lambda}}$ in $T^*\mathcal{F}_{\boldsymbol{\lambda}}$ as the zero section and consider the points x_I as points of $T^*\mathcal{F}_{\boldsymbol{\lambda}}$.

4.2. **Schubert cells, conormal bundles.** For any $\sigma \in S_n$, we consider the coordinate flag in \mathbb{C}^n ,

$$V^\sigma : 0 = V_0 \subset V_1 \subset \dots \subset V_n = \mathbb{C}^n,$$

where V_i is the span of $\epsilon_{\sigma(1)}, \dots, \epsilon_{\sigma(i)}$. For $I \in \mathcal{I}_{\boldsymbol{\lambda}}$ we define the *Schubert cell*

$$(4.1) \quad \Omega_{\sigma, I} = \{F \in \mathcal{F}_{\boldsymbol{\lambda}} \mid \dim(F_p \cap V_q^\sigma) = \#\{i \in I_1 \cup \dots \cup I_p \mid \sigma^{-1}(i) \leq q\} \forall p \leq N, \forall q \leq n\}.$$

The Schubert cell $\Omega_{\sigma, I}$ is an affine space of dimension

$$(4.2) \quad \ell_{\sigma, I} = \#\{(a, b) \in \{1, \dots, n\}^2 \mid \sigma(a) \in I_j, \sigma(b) \in I_k, a > b, j < k\}.$$

For a fixed σ , the flag manifold is the disjoint union of the cells $\Omega_{\sigma, I}$. We have $x_I \in \Omega_{\sigma, I}$, see e.g. [FP, Sect.2.2].

For $\sigma \in S_n$, recall the combinatorial partial ordering \leq_σ on \mathcal{I}_λ defined in Section 2.7. It is the same as the *geometric* partial ordering on \mathcal{I}_λ : for $I, J \in \mathcal{I}_\lambda$, we have that $J \leq_\sigma I$ if and only if x_J lies in the closure of $\Omega_{\sigma, I}$.

The Schubert cell $\Omega_{\sigma, I}$ is a smooth submanifold of \mathcal{F}_λ , hence we can consider its conormal space

$$C\Omega_{\sigma, I} = \{\alpha \in \pi^{-1}(\Omega_{\sigma, I}) \mid \alpha(T_{\pi(\alpha)}\Omega_{\sigma, I}) = 0\} \subset T^*\mathcal{F}_\lambda,$$

cf. the Remark in Section 4.6. The conormal space $C\Omega_{\sigma, I}$ is the total space of a vector subbundle of $T^*\mathcal{F}_\lambda$ over $\Omega_{\sigma, I}$. The rank of this subbundle is $\dim \mathcal{F}_\lambda - \dim \Omega_{\sigma, I}$. Hence, as a manifold $C\Omega_{\sigma, I}$ is an affine cell of dimension $\dim \mathcal{F}_\lambda$. In particular, its dimension is independent of σ, I . Define

$$(4.3) \quad \text{Slope}_{\sigma, I} = \bigcup_{J \leq_\sigma I} C\Omega_{\sigma, J}.$$

4.3. Torus action. The diagonal action of the torus $(\mathbb{C}^\times)^n$ on \mathbb{C}^n induces an action on \mathcal{F}_λ , and hence on the cotangent bundle $T^*\mathcal{F}_\lambda$. We extend this $(\mathbb{C}^\times)^n$ -action to the action of $T = (\mathbb{C}^\times)^n \times \mathbb{C}^\times$ so that the extra \mathbb{C}^\times acts on the fibers of $T^*\mathcal{F}_\lambda \rightarrow \mathcal{F}_\lambda$ by multiplication.

The fixed points of T acting on $T^*\mathcal{F}_\lambda$ are the points x_I described above.

In the next sections we consider the equivariant K -theory and equivariant elliptic cohomology of $T^*\mathcal{F}_\lambda$ with respect to this T -action.

4.4. Equivariant K -theory of $T^*\mathcal{F}_\lambda$. We consider the equivariant K -theory algebra $K_T(T^*\mathcal{F}_\lambda)$. Our general reference for equivariant K -theory is [ChG, Ch.5].

Denote $S_\lambda = S_{\lambda_1} \times \dots \times S_{\lambda_N}$ the product of symmetric groups. Consider variables $\Gamma_i = \{\gamma_{i,1}, \dots, \gamma_{i,\lambda_i}\}$, $i = 1, \dots, N$. Let $\mathbf{\Gamma} = (\Gamma_1; \dots; \Gamma_N)$. The group S_λ acts on the set $\mathbf{\Gamma}$ by permuting the variables with the same first index. Let $\mathbb{C}[\mathbf{\Gamma}^{\pm 1}]$ be the algebra of Laurent polynomials in variables $\gamma_{i,j}$ and $\mathbb{C}[\mathbf{\Gamma}^{\pm 1}]^{S_\lambda}$ the subalgebra of invariants with respect to the S_λ -action.

Consider variables $\mathbf{z} = \{z_1, \dots, z_n\}$ and h . The group S_n acts on the set \mathbf{z} by permutations. Let $\mathbb{C}[\mathbf{z}^{\pm 1}, h^{\pm 1}]$ be the algebra of Laurent polynomials in variables \mathbf{z}, h and $\mathbb{C}[\mathbf{z}^{\pm 1}, h^{\pm 1}]^{S_n}$ the subalgebra of invariants with respect to the S_n -action. We have

$$(4.4) \quad K_T(T^*\mathcal{F}_\lambda) = \mathbb{C}[\mathbf{\Gamma}^{\pm 1}]^{S_\lambda} \otimes \mathbb{C}[\mathbf{z}^{\pm 1}, h^{\pm 1}] / \langle f(\mathbf{\Gamma}) = f(\mathbf{z}) \text{ for any } f \in \mathbb{C}[\mathbf{z}^{\pm 1}]^{S_n} \rangle.$$

Here $\gamma_{i,j}$ correspond to (virtual) line bundles also denoted by $\gamma_{i,j}$ with

$$\bigoplus_{j=1}^{\lambda_i} \gamma_{i,j} = F_i / F_{i-1},$$

while z_a and h correspond to the factors of $T = (\mathbb{C}^\times)^n \times \mathbb{C}^\times$.

The algebra $K_T(T^*\mathcal{F}_\lambda)$ is a module over $K_T(pt; \mathbb{C}) = \mathbb{C}[\mathbf{z}^{\pm 1}, h^{\pm 1}]$.

4.5. Equivariant localization – moment map description. Consider the *equivariant localization* map

$$(4.5) \quad \text{Loc} : K_T(T^*\mathcal{F}_\lambda) \rightarrow K_T((T^*\mathcal{F}_\lambda)^T) = \bigoplus_{I \in \mathcal{I}_\lambda} K_T(x_I)$$

whose components are the restrictions to the fixed points x_I . Restriction of the class $\alpha \in K_T(T^*\mathcal{F}_\lambda)$ to the fix point x_I will be denoted by $\alpha|_{x_I}$. In terms of the variables $\mathbf{\Gamma}$ restriction to x_I is the substitution

$$(4.6) \quad \{\gamma_{k,1}, \dots, \gamma_{k,\lambda_k}\} \mapsto \{z_a \mid a \in I_k\} \quad \text{for all } k = 1, \dots, N.$$

Equivariant localization theory (see e.g. [ChG, Ch.5], [RoKu, Appendix]) asserts that Loc is an injection of algebras. Moreover, an element of the right-hand side is in the image of Loc if the difference of the I -th and $s_{I,J}(I)$ -th components is divisible by $1 - z_i/z_j$ in $\mathbb{C}[z^{\pm 1}, h^{\pm 1}]$ for all $I \in \mathcal{I}_\lambda$ and $i, j \in \{1, \dots, n\}$. Here $s_{I,J}(I)$ is the partition obtained from I by switching the numbers i and j .

4.6. Normal Euler classes of conormal spaces at torus fixed points. Define the following classes

$$(4.7) \quad e_{\sigma,I,+}^{\text{hor}} = \prod_{k < l} \prod_{\sigma(a) \in I_k} \prod_{\substack{\sigma(b) \in I_l \\ b < a}} (1 - z_{\sigma(a)}/z_{\sigma(b)}), \quad e_{\sigma,I,-}^{\text{hor}} = \prod_{k < l} \prod_{\sigma(a) \in I_k} \prod_{\substack{\sigma(b) \in I_l \\ b > a}} (1 - z_{\sigma(a)}/z_{\sigma(b)}),$$

$$(4.8) \quad e_{\sigma,I,+}^{\text{vert}} = \prod_{k < l} \prod_{\sigma(a) \in I_k} \prod_{\substack{\sigma(b) \in I_l \\ b > a}} (1 - h z_{\sigma(b)}/z_{\sigma(a)}), \quad e_{\sigma,I,-}^{\text{vert}} = \prod_{k < l} \prod_{\sigma(a) \in I_k} \prod_{\substack{\sigma(b) \in I_l \\ b < a}} (1 - h z_{\sigma(b)}/z_{\sigma(a)}),$$

in $K_T(x_I) = \mathbb{C}[z^{\pm 1}, h^{\pm 1}]$. We also set $e_{\sigma,I} = e_{\sigma,I,-}^{\text{hor}} e_{\sigma,I,-}^{\text{vert}}$.

Recall that if \mathbb{C}^\times acts on a line \mathbb{C} by $\alpha \cdot x = \alpha^r x$, then the \mathbb{C}^\times -equivariant Euler class of the line bundle $\mathbb{C} \rightarrow \{0\}$ is $e(\mathbb{C} \rightarrow \{0\}) = 1 - z^r \in K_{\mathbb{C}^\times}(\text{point}) = \mathbb{C}[z^{\pm 1}]$. Thus standard knowledge on the tangent bundle of flag manifolds imply that

$$(4.9) \quad e(T\Omega_{\sigma,I}|_{x_I}) = e_{\sigma,I,+}^{\text{hor}}, \quad e(\nu(\Omega_{\sigma,I} \subset \mathcal{F}_\lambda)|_{x_I}) = e_{\sigma,I,-}^{\text{hor}},$$

where $\nu(A \subset B)$ means the normal bundle of a submanifold A in the ambient manifold B , and $\xi|_x$ means the restriction of the bundle ξ over the point x in the base space. Therefore we also have

$$e(C\Omega_{\sigma,I}|_{x_I}) = e_{\sigma,I,+}^{\text{vert}}, \quad e((\pi^{-1}(\Omega_{\sigma,I}) - C\Omega_{\sigma,I})|_{x_I}) = e_{\sigma,I,-}^{\text{vert}},$$

where $C\Omega_{\sigma,I}$ and $\pi^{-1}(\Omega_{\sigma,I})$ are considered bundles over $\Omega_{\sigma,I}$. Now consider $C\Omega_{\sigma,I}$ as a(n open) submanifold of $T^*\mathcal{F}_\lambda$. Then we obtain

$$(4.10) \quad e(\nu(C\Omega_{\sigma,I} \subset T^*\mathcal{F}_\lambda)|_{x_I}) = e_{\sigma,I,-}^{\text{hor}} e_{\sigma,I,-}^{\text{vert}} = e_{\sigma,I}.$$

Remark. For a permutation $\sigma \in S_n$, let the cocharacter $\chi : \mathbb{C}^\times \rightarrow (\mathbb{C}^\times)^n$, $t \mapsto (t^{w_1}, \dots, t^{w_n})$ satisfies the property: $w_{\sigma(a)} > w_{\sigma(b)}$ for all $a > b$. The formulas of this subsection show that $C\Omega_{\sigma,I} = \{x \in T^*\mathcal{F}_\lambda \mid \lim_{t \rightarrow 0} \chi(t)x = x_I\}$. This latter set is a special case for $T^*\mathcal{F}_\lambda$ of the notion ‘‘stable leaf’’ of the fixed point x_I corresponding to the ‘‘chamber’’ $\{(w_1, \dots, w_n) \in \mathbb{R}^n \mid w_{\sigma(a)} > w_{\sigma(b)} \text{ for all } a > b\} \subset \text{Cochar}((\mathbb{C}^\times)^n) \otimes_{\mathbb{Z}} \mathbb{R} = \mathbb{R}^n$ in [MO1, Section 3.2.2]. Thus the notion of ‘‘chambers’’ of [MO1] correspond to the permutations $\sigma \in S_n$ in this paper.

5. K -THEORETIC STABLE ENVELOPES

5.1. **Definition of K -theoretic stable envelopes.** The K -theoretic stable envelope maps for Nakajima varieties X are defined by Maulik and Okounkov in [MO2]. That definition is reproduced in [O, Section 9.1] and [OS, Section 2.1]. Here we recall their axiomatic definition of stable envelope classes (images of coordinate vectors under the stable envelope map) in the special case $X = T^*\mathcal{F}_\lambda$ acted upon by $(\mathbb{C}^*)^n \times \mathbb{C}^*$.

Let $\Delta \subset \mathbb{R}^N$ be an alcove and $\nu \in \Delta$. Define the virtual ‘‘slope’’ bundle

$$S_\nu = \prod_{k=1}^N \prod_{a=1}^{\lambda_k} \gamma_{k,i}^{-\nu_k} \in \text{Pic}(T^*\mathcal{F}_\lambda) \otimes_{\mathbb{Z}} \mathbb{R}.$$

Observe that S_ν restricted to the fixed point x_I is S_I from Section 3.5.

Definition. ([MO2], [O, Section 9.1], [OS, Section 2.1]) An element $\text{Stab}_{\sigma,I}^\Delta \in K_T(T^*\mathcal{F}_\lambda)$ for an alcove $\Delta \subset \mathbb{R}^N$, $\sigma \in S_n$, $I \in \mathcal{I}_\lambda$ is called the *stable envelope class*, if it satisfies the following axioms:

- (I) $\text{Stab}_{\sigma,I}^\Delta$ is supported on $\text{Slope}_{\sigma,I}$;
- (II) $\text{Stab}_{\sigma,I}^\Delta|_{x_I} = h^{\text{codim}(\Omega_{\sigma,I} \subset \mathcal{F}_\lambda)/2} P_{\sigma,I} e_{\sigma,I}$;
- (III) $\mathcal{N}(\text{Stab}_{\sigma,I}^\Delta|_{x_I} \cdot S_\nu|_{x_I}) \subset \mathcal{N}(\text{Stab}_{\sigma,J}^\Delta|_{x_J} \cdot S_\nu|_{x_J})$, if $\nu \in \Delta$.

Observe that while condition (III) depends on $\nu \in \mathbb{R}^N$, the class $\text{Stab}_{\sigma,I}^\Delta$ only depends on the alcove ν belongs to. Namely, the inclusion in (III) can be reformulated as

$$\mathcal{N}(\text{Stab}_{\sigma,I}^\Delta|_{x_I}) + \mathcal{N}\left(\frac{S_\nu|_{x_I}}{S_\nu|_{x_J}}\right) \subset \mathcal{N}(\text{Stab}_{\sigma,J}^\Delta|_{x_J}),$$

where $+$ is the Minkowski sum and $\mathcal{N}\left(\frac{S_\nu|_{x_I}}{S_\nu|_{x_J}}\right)$ is a point. Thus (III) says that the Newton polytope of $\text{Stab}_{\sigma,I}^\Delta|_{x_I}$ lies inside the Newton polytope of $\text{Stab}_{\sigma,J}^\Delta|_{x_J}$ when shifted by the vector $\mathcal{N}\left(\frac{S_\nu|_{x_I}}{S_\nu|_{x_J}}\right)$ which depends on ν in Δ .

Remark. In fact, in [MO2, O, OS] the definition of stable envelope classes also depends on a choice of distinguishing half of the normal directions at each torus fixed point. For $T^*\mathcal{F}_\lambda$ there is a natural choice – distinguishing the cotangent directions – that we make throughout the paper. Moreover, condition (II) in the cited papers reads as follows

$$(5.1) \quad \text{Stab}_{\sigma,I}^\Delta|_{x_I} = (-1)^{\dim(\text{distinguished directions in } C\Omega_{\sigma,I})} \sqrt{\frac{\det(\nu_{\sigma,I})}{\det(T^*\mathcal{F}_\lambda|_{x_I})}} e(\nu_{\sigma,I}),$$

where $\nu_{\sigma,I}$ is the normal bundle of $C\Omega_{\sigma,I}$ at x_I . However, using the notation

$$\begin{aligned} d_{\sigma,I,+}^{\text{hor}} &= \prod_{k < l} \prod_{\sigma(a) \in I_k} \prod_{\substack{\sigma(b) \in I_l \\ b < a}} \frac{z_{\sigma(b)}}{z_{\sigma(a)}}, & d_{\sigma,I,-}^{\text{hor}} &= \prod_{k < l} \prod_{\sigma(a) \in I_k} \prod_{\substack{\sigma(b) \in I_l \\ b > a}} \frac{z_{\sigma(b)}}{z_{\sigma(a)}}, \\ d_{\sigma,I,+}^{\text{vert}} &= \prod_{k < l} \prod_{\sigma(a) \in I_k} \prod_{\substack{\sigma(b) \in I_l \\ b > a}} \frac{z_{\sigma(a)}}{h z_{\sigma(b)}}, & d_{\sigma,I,-}^{\text{vert}} &= \prod_{k < l} \prod_{\sigma(a) \in I_k} \prod_{\substack{\sigma(b) \in I_l \\ b < a}} \frac{z_{\sigma(a)}}{h z_{\sigma(b)}} \end{aligned}$$

the right hand side of equation (5.1) is

$$\prod_{k < l} \prod_{\sigma(a) \in I_k} \prod_{\substack{\sigma(b) \in I_l \\ b > a}} (-1) \cdot \sqrt{\frac{d_{\sigma, I, -}^{\text{hor}} d_{\sigma, I, -}^{\text{vert}}}{d_{\sigma, I, +}^{\text{vert}} d_{\sigma, I, -}^{\text{vert}}}} \cdot e_{\sigma, I} =$$

$$\prod_{k < l} \prod_{\sigma(a) \in I_k} \prod_{\substack{\sigma(b) \in I_l \\ b > a}} \left(-\frac{z_{\sigma(b)}}{z_{\sigma(a)}} \right) h^{\text{codim}(\Omega_{\sigma, I} \subset \mathcal{F}_\lambda)/2} e_{\sigma, I} = h^{\text{codim}(\Omega_{\sigma, I} \subset \mathcal{F}_\lambda)/2} P_{\sigma, I} e_{\sigma, I},$$

as stated in the definition above.

It is proved in [O, Theorem 9.2.1] that stable envelope classes – if exist – are unique. Their existence statement for Nakajima varieties is announced to be published in [MO2], see [O, 9.1.1], cf. also Theorem 2 and Proposition 3.5 in [AO].

5.2. Weight functions project to stable envelope classes. We substitute the variables

$$t_1^{(k)}, \dots, t_{\lambda^{(k)}}^{(k)} \mapsto \gamma_{1,1}, \dots, \gamma_{1,\lambda_1}, \quad \gamma_{2,1}, \dots, \gamma_{2,\lambda_2}, \quad \dots, \quad \gamma_{k,1}, \dots, \gamma_{k,\lambda_k}$$

into the function $\tilde{W}_{\sigma, I}^\Delta(\mathbf{t}, \mathbf{z}, h)$ (see Section 3). The resulting function will be denoted by $\tilde{W}_{\sigma, I}^\Delta(\mathbf{\Gamma}, \mathbf{z}, h)$. Observe that we have

$$\tilde{W}_{\sigma, I}^\Delta(\mathbf{\Gamma}, \mathbf{z}, h)|_{x_J} = \tilde{W}_{\sigma, I}^\Delta(z_J, \mathbf{z}, h).$$

In view of Lemma 3.3 this implies that the function $\tilde{W}_{\sigma, I}^\Delta(\mathbf{\Gamma}, \mathbf{z}, h)$ – although not a Laurent polynomial itself – has the property that all its x_J restrictions are Laurent polynomials. The restrictions also satisfy the divisibility property of Section 4.5, hence equivariant localization theory implies that there is a unique element in $K_T(T^*\mathcal{F}_\lambda)$ whose x_J restrictions are the same as those of $\tilde{W}_{\sigma, I}^\Delta(\mathbf{\Gamma}, \mathbf{z}, h)$. By slight abuse of language we will denote this element of $K_T(T^*\mathcal{F}_\lambda)$ by $[\tilde{W}_{\sigma, I}^\Delta(\mathbf{\Gamma}, \mathbf{z}, h)]$.

Theorem 5.1. *Stable envelope classes $\text{Stab}_{\sigma, I}^\Delta$ exist for $T^*\mathcal{F}_\lambda$, namely,*

$$(5.2) \quad \text{Stab}_{\sigma, I}^\Delta = h^{\text{codim}(\Omega_{\sigma, I} \subset \mathcal{F}_\lambda)/2} \cdot [\tilde{W}_{\sigma, I}^\Delta(\mathbf{\Gamma}, \mathbf{z}, h)].$$

Proof. We denote the right hand side of (5.2) by $\omega_{\sigma, I}$ and we will prove that it satisfies axioms (I)-(III).

Axiom (II) follows from Lemma 3.5. To prove axiom (III) observe that $S_\nu|_{x_I} = S_I$ and hence Theorem 3.9 claims

$$\mathcal{N}(W_I^\Delta(\mathbf{z}_J, \mathbf{z}, h) S_\nu|_{x_I}) \subset \mathcal{N}(W_J^\Delta(\mathbf{z}_J, \mathbf{z}, h) S_\nu|_{x_J}).$$

Writing this statement for the pair $\sigma^{-1}(I)$, $\sigma^{-1}(J)$ instead of I, J , we obtain

$$\mathcal{N}(W_{\sigma^{-1}(I)}^\Delta(\mathbf{z}_{\sigma^{-1}(J)}, \mathbf{z}, h) S_\nu|_{x_{\sigma^{-1}(I)}}) \subset \mathcal{N}(W_{\sigma^{-1}(J)}^\Delta(\mathbf{z}_{\sigma^{-1}(J)}, \mathbf{z}, h) S_\nu|_{x_{\sigma^{-1}(J)}}).$$

Replace z_a with $z_{\sigma(a)}$ for $a = 1, \dots, n$, to get

$$\mathcal{N}(W_{\sigma^{-1}(I)}^\Delta(\mathbf{z}_J, \mathbf{z}_\sigma, h) S_\nu|_{x_I}) \subset \mathcal{N}(W_{\sigma^{-1}(J)}^\Delta(\mathbf{z}_J, \mathbf{z}_\sigma, h) S_\nu|_{x_J}),$$

where $\mathbf{z}_\sigma = (z_{\sigma(1)}, \dots, z_{\sigma(n)})$. Using the definition of $W_{\sigma, I}$ functions this translates to

$$\mathcal{N}(W_{\sigma, I}^\Delta(\mathbf{z}_J, \mathbf{z}, h) S_\nu|_{x_I}) \subset \mathcal{N}(W_{\sigma, J}^\Delta(\mathbf{z}_J, \mathbf{z}, h) S_\nu|_{x_J}).$$

Since the ratio between $W_I^\Delta(\mathbf{z}_J, \mathbf{z}, h)$ and $\tilde{W}_I^\Delta(\mathbf{z}_J, \mathbf{z}, h)$ is a universal expression (not depending on I) the cancellation property of Minkowski sums gives

$$\mathcal{N}(\tilde{W}_{\sigma, I}^\Delta(\mathbf{z}_J, \mathbf{z}, h)S_\nu|_{x_I}) \subset \mathcal{N}(\tilde{W}_{\sigma, J}^\Delta(\mathbf{z}_J, \mathbf{z}, h)S_\nu|_{x_J}),$$

what we wanted to prove.

The rest of this section is the proof of axiom (I). In the proof we will consider restrictions of elements of $K_T(T^*\mathcal{F}_\lambda)$ to x_J , to $\Omega_{\sigma, J}$, and to $\pi^{-1}(\Omega_{\sigma, J})$. Observe that algebraically these three restrictions are the same maps, because the last two sets are topological cells, equivariantly homotopy equivalent to the one-point set x_J .

Lemma 5.2. *Let \prec be a linear order refining \leq_σ , and let $J^{(1)} \prec J^{(2)} \prec \dots \prec J^{(d)}$ be the elements of \mathcal{I}_λ , that is $d = |\mathcal{I}_\lambda|$. Let $a \in \{2, \dots, d\}$. If a class $\omega \in K_T(T^*\mathcal{F}_\lambda)$ restricted to*

$$\cup_{i=a}^d \pi^{-1}(\Omega_{\sigma, J^{(i)}})$$

is 0, and $\omega|_{x_{J^{(a-1)}}} = 0$, then ω restricted to

$$\cup_{i=a-1}^d \pi^{-1}(\Omega_{\sigma, J^{(i)}})$$

is also 0.

Proof. This is a well known argument involving a Mayer-Vietoris and a Gysin sequence, and depending on the non-vanishing of a normal Euler class (which holds for partial flag varieties). See for example [FR, Lemma 3.6] for a detailed argument in cohomology. \square

Lemma 5.3. *Let T act on the vector space X with an invariant subspace Y . Let e be the T -equivariant normal Euler class of $Y \subset X$. Then the following conditions are equivalent for a class in $K_T(X)$:*

- *it is supported on Y ;*
- *it is divisible by e .*

Proof. The statement follows from the exactness of the Gysin sequence

$$K_T(Y) \xrightarrow{e} K_T(Y) \longrightarrow K_T(X - Y)$$

where the second map is the composition of the isomorphism $K_T(Y) = K_T(X)$ and the restriction to $X - Y$. \square

Lemma 5.4. *We have*

$$\omega_{\sigma, I} = \sum_{J \leq_\sigma I} f_{IJ}(\mathbf{z}, h) [\overline{C\Omega_{\sigma, J}}]$$

for some Laurent polynomials $f_{IJ}(\mathbf{z}, h)$.

Proof. Let \prec be a total order refining \leq_σ , and let $J^{(1)} \prec J^{(2)} \prec \dots \prec J^{(r)} = I$ be the elements of \mathcal{I}_λ that are $\leq_\sigma I$. Let $k \in \{0, \dots, r\}$. We will prove the following statement by induction on k .

(*k) *There exist Laurent polynomials $f_{IJ^{(i)}}(\mathbf{z}, h)$ for $i \in \{r - k + 1, \dots, r\}$ such that*

$$(5.3) \quad \omega_{\sigma, I} - \sum_{i=r-k+1}^r f_{IJ^{(i)}}(\mathbf{z}, h) [\overline{C\Omega_{\sigma, J^{(i)}}}]$$

is supported on $\cup_{i=1}^{r-k} \pi^{-1}(\Omega_{\sigma, J^{(i)}})$.

Statement (*k) for $k = 0$ follows from the iterated application of Lemma 5.2, using Lemma 3.4.

For the induction step $k \mapsto k+1$, consider (5.3) restricted to $x_{J^{(r-k)}}$. The term $\omega_{\sigma, I}|_{x_{J^{(r-k)}}$ is divisible by $e_{\sigma, J^{(r-k)}, -}^{\text{vert}}$ by Lemma 3.6. The classes $[\overline{C\Omega_{\sigma, J^{(i)}}}]$ appearing in (5.3), restricted to $x_{J^{(r-k)}}$ are also divisible by $e_{\sigma, J^{(r-k)}, -}^{\text{vert}}$, due to the fact that the Schubert cell stratification is a Whitney stratification. Therefore (5.3), restricted to $x_{J^{(r-k)}}$ is divisible by $e_{\sigma, J^{(r-k)}, -}^{\text{vert}}$.

We claim that (5.3), restricted to $x_{J^{(r-k)}}$ is also divisible by $e_{\sigma, J^{(r-k)}, -}^{\text{hor}}$. This follows from the induction hypotheses and Lemma 5.3. We obtained that (5.3) restricted to $x_{J^{(r-k)}}$ can be written as $f_{IJ^{(r-k)}}(\mathbf{z}, h) \cdot e_{\sigma, J^{(r-k)}, -}^{\text{hor}} e_{\sigma, J^{(r-k)}, -}^{\text{vert}}$ for some Laurent polynomial $f_{IJ^{(r-k)}}(\mathbf{z}, h)$. Therefore the class

$$\left(\omega_{\sigma, I} - \sum_{i=r-k-1}^r f_{IJ^{(i)}}(\mathbf{z}, h) [\overline{C\Omega_{\sigma, J^{(i)}}}] \right) - f_{IJ^{(r-k)}}(\mathbf{z}, h) [\overline{C\Omega_{\sigma, J^{(r-k)}}}]$$

restricted to $x_{J^{(r-k)}}$ is 0. Since it is also 0 when restricted to $\cup_{i=r-k-1}^r \pi^{-1}(\Omega_{\sigma, J^{(i)}})$ (by induction and the property of the order \prec), Lemma 5.3 gives the induction step.

Statement (*k) for $k = r$ proves Lemma 5.4. \square

The class $[\overline{C\Omega_{\sigma, I}}]$ is supported on $\text{Slope}_{\sigma, I}$, and clearly $J \leq_{\sigma} I$ implies $\text{Slope}_{\sigma, J} \subset \text{Slope}_{\sigma, I}$, hence Lemma 5.4 proves that $\omega_{\sigma, I}$ satisfies axiom (III). This finishes the proof of Theorem 5.1. \square

Remark. Our arguments show that in the definition of $\text{Stab}_{\sigma, I}$ axiom (I) can be replaced with its local version

(I') $\text{Stab}_{\sigma, I}^{\Delta}|_{x_J}$ is divisible by $e_{\sigma, J, -}^{\text{vert}}$ as was done, for example, in [RTV2, Theorem 3.1].

Remark. The functions $f_{IJ}(\mathbf{z}, h)$ appearing in Lemma 5.4 are related to the notion of “local Euler obstruction” in cohomology.

5.3. Note on R -matrices. A key feature of the $\text{Stab}_{\sigma, I}^{\Delta}$ classes is that some endomorphisms encoded by them satisfy Yang-Baxter equations, that is, they are R -matrices. For more details, see eg. [O, RTV2]. For completeness here we show the prototype R -matrices for the convention used in this paper.

Let $\mathcal{X}_n = \cup_{|\lambda|=n} T^* \mathcal{F}_{\lambda}$. For $I \in \mathcal{I}_{\lambda}$ let $v_I = v_{i_1} \otimes \dots \otimes v_{i_n} \in (\mathbb{C}^N)^{\otimes n}$ for $i_j = k$ if $j \in I_k$. Stable envelope maps are defined

$$\text{Stab}_{\sigma}^{\Delta} : (\mathbb{C}^N)^{\otimes n} \otimes \mathbb{C}(\mathbf{z}, h) = K_T(\mathcal{X}_n^T) \otimes \mathbb{C}(\mathbf{z}, h) \rightarrow K_T(\mathcal{X}_n) \otimes \mathbb{C}(\mathbf{z}, h)$$

by $v_I \mapsto \text{Stab}_{\sigma, I}^{\Delta}$.

Stable envelope maps define “geometric” R -matrices by

$$\mathcal{R}_{\sigma', \sigma}^{\Delta} = (\text{Stab}_{\sigma'}^{\Delta})^{-1} \circ \text{Stab}_{\sigma}^{\Delta}.$$

For example, for $n = 2, N = 2$, we have

$$\begin{aligned}\text{Stab}_{\text{id}}^{\Delta}(v_1 \otimes v_2) &= (\gamma/z_1)^{m_{1,2}} (1 - z_2/\gamma) \cdot \sqrt{h} \\ \text{Stab}_{\text{id}}^{\Delta}(v_2 \otimes v_1) &= (\gamma/z_2)^{m_{1,2}} (1 - h z_1/\gamma) \\ \text{Stab}_s^{\Delta}(v_1 \otimes v_2) &= (\gamma/z_1)^{m_{1,2}} (1 - h z_2/\gamma) \\ \text{Stab}_s^{\Delta}(v_2 \otimes v_1) &= (\gamma/z_2)^{m_{1,2}} (1 - z_1/\gamma) \cdot \sqrt{h}.\end{aligned}$$

Therefore, $\mathcal{R}_{s,\text{id}}^{\Delta}$ acting on $(\mathbb{C}^2)^{\otimes 2}$ with ordered basis $v_1 \otimes v_1, v_1 \otimes v_2, v_2 \otimes v_1, v_2 \otimes v_2$ is

$$(1) \oplus \left(\begin{array}{cc} \frac{\sqrt{h}(1 - z_2/z_1)}{1 - h z_2/z_1} & \frac{(1 - h)(z_2/z_1)^{-m_{1,2}}}{1 - h z_2/z_1} \\ \frac{(1 - h)(z_2/z_1)^{m_{1,2}+1}}{1 - h z_2/z_1} & \frac{\sqrt{h}(1 - z_2/z_1)}{1 - h z_2/z_1} \end{array} \right) \oplus (1).$$

The R -matrix $R_{s,\text{id}}$ used in [RTV2]

$$(1) \oplus \left(\begin{array}{cc} \frac{1 - z_2/z_1}{1 - h z_2/z_1} & \frac{(1 - h)}{1 - h z_2/z_1} \\ \frac{(1 - h)z_2/z_1}{1 - h z_2/z_1} & \frac{h(1 - z_2/z_1)}{1 - h z_2/z_1} \end{array} \right) \oplus (1)$$

is obtained from this one by putting $m_{1,2} = 0$ (considering the anti-dominant alcove) and a change in the normalization convention involving h .

Similar calculation shows that for general N , but $n = 2$, the operator $\mathcal{R} = \mathcal{R}_{s,\text{id}}^{\Delta}$ acting on $(\mathbb{C}^N)^{\otimes 2}$ has entries

$$(5.4) \quad \mathcal{R}_{jj}^{jj} = 1, \quad \mathcal{R}_{jk}^{jk} = \sqrt{h} \frac{1 - z_2/z_1}{1 - h z_2/z_1} \quad (1 \leq j, k \leq N, j \neq k),$$

$$\mathcal{R}_{kj}^{jk} = \frac{(1 - h)(z_2/z_1)^{m_{j,k}+1}}{1 - h z_2/z_1}, \quad \mathcal{R}_{jk}^{kj} = \frac{(1 - h)(z_2/z_1)^{-m_{j,k}}}{1 - h z_2/z_1} \quad (1 \leq j < k \leq N),$$

and otherwise 0, while the R -matrix $R = R_{s,\text{id}}$ used in [RTV2] has entries $R_{jj}^{jj} = 1$,

$$(5.5) \quad R_{jk}^{jk} = \frac{1 - z_2/z_1}{1 - h z_2/z_1} \quad (j < k), \quad R_{jk}^{jk} = h \frac{1 - z_2/z_1}{1 - h z_2/z_1} \quad (k < j),$$

$$R_{kj}^{jk} = \frac{(1 - h)(z_2/z_1)}{1 - h z_2/z_1}, \quad R_{jk}^{kj} = \frac{1 - h}{1 - h z_2/z_1} \quad (1 \leq j < k \leq N),$$

and otherwise 0.

6. LINE BUNDLES AND QUADRATIC FORMS

6.1. Line bundles on \mathbb{C}^p . Let $E = \mathbb{C}/\Lambda$ be an elliptic curve, where $\Lambda = \mathbb{Z} + \tau\mathbb{Z}$, $\tau \in \mathbb{C}$, $\text{Im } \tau > 0$, so that $E^p = \mathbb{C}^p/\Lambda^p$. For each pair (M, v) consisting of a symmetric integral $p \times p$ matrix M and $v \in (\mathbb{R}/\mathbb{Z})^p$ let $\mathcal{L}(M, v)$ be the line bundle $(\mathbb{C}^p \times \mathbb{C})/\Lambda^p \rightarrow \mathbb{C}^p/\Lambda^p$ with action

$$\lambda \cdot (x, u) = (x + \lambda, e_{\lambda}(x)(u)), \quad \lambda \in \Lambda^p, x \in \mathbb{C}^p, u \in \mathbb{C},$$

and cocycle

$$e_{n+m\tau}(x) = (-1)^{n^t M n} (-e^{i\pi\tau})^{m^t M m} e^{2\pi i m^t (Mx+v)}, \quad n, m \in \mathbb{Z}^p.$$

To an integral symmetric $p \times p$ matrix M and a vector $v \in \mathbb{C}^p$ we associate the integral quadratic form $M(x) = x^t M x$ and the linear form $v(x) = x^t v$ on the universal cover \mathbb{C}^p of E^p , and we call them the quadratic form and the linear form of the line bundle $\mathcal{L}(M, v)$. The linear form is defined up to addition of an integral linear form.

Proposition 6.1 ([FRV, Proposition 5.1], see [Mu, Section I.2]).

- (i) $\mathcal{L}(M, v)$ is isomorphic to $\mathcal{L}(M', v')$ if and only if $M = M'$ and $v \equiv v' \pmod{\Lambda^p}$.
- (ii) For generic E , every holomorphic line bundle on E^p is isomorphic to $\mathcal{L}(M, v)$ for some (M, v) .
- (iii) $\mathcal{L}(M_1, v_1) \otimes \mathcal{L}(M_2, v_2) \cong \mathcal{L}(M_1 + M_2, v_1 + v_2)$.
- (iv) Let $\sigma \in S_p$ act by permutations on E^p and \mathbb{C}^p . Denote also by σ the corresponding $p \times p$ permutation matrix. Then

$$\sigma^* \mathcal{L}(M, v) = \mathcal{L}(\sigma^t M \sigma, \sigma^t v).$$

Remark. Sections of $\mathcal{L}(M, v)$ are the same as functions f on \mathbb{C}^p such that $f(x + \lambda) = e_{\lambda}(x)^{-1} f(x)$ for all $\lambda \in \Lambda^p$, $x \in \mathbb{C}^p$. Explicitly, a function on \mathbb{C}^p defines a section of $\mathcal{L}(M, v)$ if and only if

$$\begin{aligned} f(x_1, \dots, x_j + 1, \dots, x_p) &= (-1)^{M_{jj}} f(x), \\ f(x_1, \dots, x_j + \tau, \dots, x_p) &= (-1)^{M_{jj}} e^{-2\pi i (\sum_k M_{jk} x_k + v_j) - \pi i \tau M_{jj}} f(x), \end{aligned}$$

for all $x \in \mathbb{C}^p$, $j = 1, \dots, p$.

Remark ([FRV]). Let $\theta(x)$ be the theta function in one variable defined by (2.3). Then, for any $r \in \mathbb{Z}^p$ and $z \in \mathbb{C}$,

$$\theta(r^t x + z) = \theta(r_1 x_1 + \dots + r_p x_p + z)$$

is a holomorphic section of $\mathcal{L}(M, v)$ with quadratic form $M(x) = (\sum_{i=1}^p r_i x_i)^2$, and linear form $v(x) = z \sum_{i=1}^p r_i x_i$. Since an integral quadratic form is an integral linear combination of squares of integral linear forms, a line bundle $\mathcal{L}(M, 0)$ has a meromorphic section which is a ratio of products of theta functions $\theta(r^t x)$ with $r \in \mathbb{Z}^p$.

6.2. Theta functions as sections of line bundles. Introduce new variables $\mathbf{v} = (v^{(1)}, \dots, v^{(N)})$, $v^{(k)} = (v_{k,1}, \dots, v_{k,\lambda_k})$, \mathbf{y} , $\boldsymbol{\nu} = (\nu_1, \dots, \nu_{N-1})$. Make the substitution

$$(6.1) \quad \begin{aligned} q, h, \mu_1/\mu_2, \dots, \mu_{N-1}/\mu_N &\mapsto e^{2\pi i \tau}, e^{2\pi i y}, e^{2\pi i \nu_1}, \dots, e^{2\pi i \nu_{N-1}}, \\ t_1^{(k)}, \dots, t_{\lambda(k)}^{(k)} &\mapsto e^{2\pi i v_{1,1}}, \dots, e^{2\pi i v_{1,\lambda_1}}, e^{2\pi i v_{2,1}}, \dots, e^{2\pi i v_{2,\lambda_2}}, \dots, e^{2\pi i v_{k,1}}, \dots, e^{2\pi i v_{k,\lambda_k}} \end{aligned}$$

for $k = 1, \dots, N$ in the functions $G_{\lambda}(\mathbf{t}, h, \boldsymbol{\mu})$, $E_{\lambda}^{\text{ell}}(\mathbf{t}, h)$. Denote the resulting functions by $\hat{G}_{\lambda}(\mathbf{v}, \mathbf{y}, \boldsymbol{\nu})$, $\hat{E}_{\lambda}^{\text{ell}}(\mathbf{v}, \mathbf{y})$.

Lemma 6.2. *The function $\hat{G}_\lambda(\mathbf{v}, y, \boldsymbol{\nu})/\hat{E}_\lambda^{\text{ell}}(\mathbf{v}, y)$ defines a meromorphic section of the line bundle $\mathcal{L}(M_\lambda, 0)$ on $E^{\lambda_1} \times \dots \times E^{\lambda_N} \times E \times E^{N-1}$, where*

$$(6.2) \quad M_\lambda(\mathbf{v}, y, \boldsymbol{\nu}) = \sum_{1 \leq j < k \leq N} \sum_{a=1}^{\lambda_j} \sum_{b=1}^{\lambda_k} (v_{j,a} - v_{k,b})^2 - y^2 \sum_{k=1}^{N-1} (\lambda_1 + \dots + \lambda_k)^2 \\ + 2 \sum_{k=1}^{N-1} (\nu_k - \lambda_k y) \sum_{j=1}^k \sum_{a=1}^{\lambda_j} v_{j,a}.$$

Proof. By straightforward calculation. □

Make the substitution

$$(6.3) \quad q, h \mapsto e^{2\pi i \tau}, e^{2\pi i y}, \\ z_1, \dots, z_n, \mu_1/\mu_2, \dots, \mu_{N-1}/\mu_N \mapsto e^{2\pi i x_1}, \dots, e^{2\pi i x_n}, e^{2\pi i \nu_1}, \dots, e^{2\pi i \nu_{N-1}},$$

in the function $G_I(\mathbf{z}, h, \boldsymbol{\mu})$. The resulting function denote by $\hat{G}_I(\mathbf{x}, y, \boldsymbol{\nu})$.

Lemma 6.3. *For $I \in \mathcal{I}_\lambda$ the function $\hat{G}_I(\mathbf{x}, y, \boldsymbol{\nu})$ defines a meromorphic section of the line bundle $\mathcal{L}(M_I, 0)$ on $E^n \times E \times E^{N-1}$, where*

$$(6.4) \quad M_I(\mathbf{x}, y, \boldsymbol{\nu}) = -2 \sum_{k=1}^{N-1} (\nu_k + \dots + \nu_{N-1}) \sum_{a \in I_k} x_a + y^2 d_I \\ + 2y \sum_{k=1}^N \sum_{a \in I_k} x_a (p_{I,k}(a) - a + 1 + n - \lambda_N).$$

Proof. By straightforward calculation. □

7. EQUIVARIANT ELLIPTIC COHOMOLOGY OF $T^*\mathcal{F}_\lambda$

7.1. Tautological bundles and Chern classes. We follow the exposition of equivariant elliptic cohomology in [FRV, Section 4], which is based on [Gr, GKV].

Let $E = \mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$, $\text{Im } \tau > 0$, be an elliptic curve and G a compact group. The G -equivariant elliptic cohomology $E_G(-)$ is a covariant functor from finite G -CW complexes to superschemes satisfying a set of axioms. For example,

$$E_{U(1)}(\text{pt}) = E, \quad E_{U(n)}(\text{pt}) = E^{(n)} = E^n/S_n.$$

For a G -space M we have the structure map $p_G : E_G(M) \rightarrow E_G(\text{pt})$. The $A = U(1)^n$ -equivariant elliptic cohomology of \mathcal{F}_λ is the fiber product obtained from the Cartesian square

$$\begin{array}{ccc} E_A(\mathcal{F}_\lambda) & \xrightarrow{x} & E^{(\lambda_1)} \times E^{(\lambda_2)} \times \dots \times E^{(\lambda_N)} \\ p_A \downarrow & & \downarrow p_\lambda \\ E^n & \xrightarrow{p_n} & E^{(n)} \end{array}$$

where the left vertical arrow is the structure map to $E_A(\text{pt})$; p_n, p_λ are canonical projections, and the top horizontal map χ is the characteristic map coming from the N equivariant bundles F_k/F_{k-1} over \mathcal{F}_λ , see [GKV], and we have

$$(7.1) \quad E_A(\mathcal{F}_\lambda) = E^n \times_{E^{(n)}} (E^{(\lambda_1)} \times E^{(\lambda_2)} \times \dots \times E^{(\lambda_N)})$$

7.2. Equivariant localization – moment map description. The inclusion of the fixed point x_I to \mathcal{F}_λ induces a map $\iota_I : E_A(\text{pt}) \rightarrow E_A(\mathcal{F}_\lambda)$ and $E_A(\mathcal{F}_\lambda)$ is the union of $\iota_I E_A(\text{pt}) \cong E^n$ where $I \in \mathcal{I}_\lambda$. The intersections of $\iota_I E_A(\text{pt}) \cong E^n$'s are described as follows. Assume $a \in I_k, b \in I_l$ for an $I \in \mathcal{I}_\lambda$, and assume $k \neq l$. Let $s_{a,b}(I)$ be obtained by switching the numbers a and b , i.e. $a \in s_{a,b}(I)_l$ and $b \in s_{a,b}(I)_k$. The “diagonal” $\Delta_{I,s_{a,b}(I)} = \{\mathbf{x} \in E^n : x_a = x_b\}$ is naturally included in both $\iota_I E_A(\text{pt}) = E^n$ and $\iota_{s_{a,b}(I)} E_A(\text{pt}) = E^n$. The counterpart of the standard GKM theory for equivariant elliptic cohomology is

$$E_A(\mathcal{F}_\lambda) = \sqcup_{I \in \mathcal{I}_\lambda} \iota_I E_A(\text{pt}) / \sim$$

where \sim is induced by gluing $\iota_I E_A(\text{pt}) = E^n$ and $\iota_{s_{a,b}(I)} E_A(\text{pt}) = E^n$ along the diagonal $\Delta_{I,s_{a,b}(I)} = \{\mathbf{x} \in E^n : x_a = x_b\}$, for any I, a, b , cf. [GKV], [GKM], [Ga, Example 4.4]. The isomorphism between the two descriptions of $E_A(\mathcal{F}_\lambda)$ is induced by the map

$$\sqcup_{I \in \mathcal{I}_\lambda} \iota_I (E_A(\text{pt})) \rightarrow E^n \times_{E^{(n)}} (E^{(\lambda_1)} \times E^{(\lambda_2)} \times \dots \times E^{(\lambda_N)})$$

whose restrictions to the copy E^n labeled by I is $\mathbf{x} \mapsto (\mathbf{x}, \mathbf{x}_{I_1}, \dots, \mathbf{x}_{I_N})$.

7.3. Cotangent bundle and the dynamical parameters. Denote $T = A \times U(1) = U(1)^n \times U(1)$, and the extra $U(1)$ acts on the fibers of $T^*\mathcal{F}_\lambda$ by multiplication. Since $T^*\mathcal{F}_\lambda$ is equivariantly homotopy equivalent to \mathcal{F}_λ we have

$$E_T(T^*\mathcal{F}_\lambda) = E_A(\mathcal{F}_\lambda) \times E,$$

a scheme over $E_T(\text{pt}) = E^n \times E$.

As in [AO] and [FRV], we consider an extended version of elliptic cohomology to accommodate for dynamical parameters in quantum group theory (or Kähler parameters in the terminology of [AO]), namely

$$\widehat{E}_T(T^*\mathcal{F}_\lambda) := E_T(T^*\mathcal{F}_\lambda) \times (\text{Pic}(T^*\mathcal{F}_\lambda) \otimes_{\mathbb{Z}} E) \simeq E_T(T^*\mathcal{F}_\lambda) \times E^{N-1},$$

a scheme over $\widehat{E}(\text{pt}) = E^{n+1+N-1}$, where the factors of E^{N-1} correspond to the line bundles $\det(F_{p+1}/F_p)$, $p = 1, \dots, N-1$ over $T^*\mathcal{F}_\lambda$.

We will denote also by ι_I the map $\widehat{E}_T(\text{pt}) \rightarrow \widehat{E}_T(T^*\mathcal{F}_\lambda)$ induced by the inclusion of the fixed point x_I into $T^*\mathcal{F}_\lambda$. Then $\widehat{E}_T(T^*\mathcal{F}_\lambda)$ consists of the components $\iota_I \widehat{E}_T(\text{pt})$ where I runs over the elements of the set \mathcal{I}_λ . By Section 7.1, we have a description of $\widehat{E}_T(T^*\mathcal{F}_\lambda)$ as a fiber product

$$(7.2) \quad \widehat{E}_T(T^*\mathcal{F}_\lambda) = (E^n \times_{E^{(n)}} (E^{(\lambda_1)} \times E^{(\lambda_2)} \times \dots \times E^{(\lambda_N)})) \times E \times E^{N-1}.$$

In particular we have the characteristic embedding

$$(7.3) \quad c : \widehat{E}_T(T^*\mathcal{F}_\lambda) \rightarrow E^n \times E^{(\lambda_1)} \times E^{(\lambda_2)} \times \dots \times E^{(\lambda_N)} \times E \times E^{N-1}$$

of the extended T -equivariant cohomology scheme into a nonsingular projective variety.

7.4. Admissible line bundles, and their sections. We will consider sections of certain “admissible” line bundles over $\widehat{E}_T(T^*\mathcal{F}_\lambda)$. These line bundles are, up to a twist by a fixed line bundle, those coming from the base scheme $\widehat{E}_T(\text{pt})$. Let p_T be the structure map

$$(7.4) \quad p_T: \widehat{E}_T(T^*\mathcal{F}_\lambda) \rightarrow \widehat{E}_T(\text{pt})$$

and $\hat{\chi} = \chi \times \text{id} \times \text{id}$ the characteristic map

$$(7.5) \quad \hat{\chi} : \widehat{E}_T(T^*\mathcal{F}_\lambda) \rightarrow E^{(\lambda_1)} \times E^{(\lambda_2)} \times \dots \times E^{(\lambda_N)} \times E \times E^{N-1}.$$

Let $\mathbf{v} = (v^{(1)}, \dots, v^{(N)})$, $v^{(k)} = (v_{k,1}, \dots, v_{k,\lambda_k})$ be coordinates on the universal cover of $E^{\lambda_1} \times \dots \times E^{\lambda_N}$ and let $y, \boldsymbol{\nu} = (\nu_1, \dots, \nu_{N-1})$ be coordinates on the universal cover of $E \times E^{N-1}$.

Let M_λ be the quadratic form defined in (6.2). Clearly M_λ is symmetric under the action of the product $S_\lambda = S_{\lambda_1} \times \dots \times S_{\lambda_N}$ of symmetric groups permuting the second index of $v_{l,a}$'s while the first index is fixed. Thus the line bundle $\mathcal{L}(M_\lambda, 0)$ on $E^{\lambda_1} \times \dots \times E^{\lambda_N} \times E \times E^{N-1}$ can be considered as a line bundle on $E^{(\lambda_1)} \times \dots \times E^{(\lambda_N)} \times E \times E^{N-1}$.

Definition. The *twisting line bundle* on $\widehat{E}_T(T^*\mathcal{F}_\lambda)$ is $\mathcal{T}_\lambda = \hat{\chi}^* \mathcal{L}(M_\lambda, 0)$.

Definition. An *admissible line bundle* on $\widehat{E}_T(T^*\mathcal{F}_\lambda)$ is a line bundle of the form

$$p_T^* \mathcal{L} \otimes \mathcal{T}_\lambda,$$

for some line bundle \mathcal{L} on $\widehat{E}_T(\text{pt})$.

We say that a meromorphic section on a complex manifold *restricts to a meromorphic section* on a submanifold if it is defined at its generic point, i.e., if the divisor of poles does not contain a component of the submanifold.

We will consider meromorphic sections of line bundles on elliptic cohomology schemes. Recall that $\widehat{E}_T(T^*\mathcal{F}_\lambda)$ has components $Y_I = \iota_I \widehat{E}_T(\text{pt})$, corresponding to the inclusion of the fixed points $x_I, I \in \mathcal{I}_\lambda$ into $T^*\mathcal{F}_\lambda$.

Definition. Let \mathcal{L} be a line bundle on $\widehat{E}_T(T^*\mathcal{F}_\lambda)$. A *meromorphic section* of \mathcal{L} is a collection of meromorphic sections s_I of $\mathcal{L}|_{Y_I}$, labeled by $I \in \mathcal{I}_\lambda$ and restricting to meromorphic sections on all intersections $Y_{I_1} \cap \dots \cap Y_{I_s}$ and such that

$$s_I|_{Y_I \cap Y_J} = s_J|_{Y_I \cap Y_J},$$

for all I, J . A *holomorphic section* is a meromorphic section whose restriction to each Y_I is holomorphic.

7.5. T -equivariant elliptic cohomology classes.

Definition. Let \mathcal{L} be a line bundle on $\widehat{E}_T(\text{pt})$. A *T -equivariant elliptic cohomology class* on $T^*\mathcal{F}_\lambda$ of degree \mathcal{L} is a holomorphic section of the admissible line bundle $p_T^* \mathcal{L} \otimes \mathcal{T}_\lambda$ on $\widehat{E}_T(T^*\mathcal{F}_\lambda)$. We denote by $H_T^{\text{ell}}(T^*\mathcal{F}_\lambda)_\mathcal{L}$ the complex vector space of T -equivariant elliptic cohomology classes of degree \mathcal{L} on $T^*\mathcal{F}_\lambda$.

7.6. Weight functions project to equivariant elliptic cohomology classes. In this section we show that the substitutions of weight functions $\tilde{W}_{\sigma,I}^{\text{ell}}$ define T -equivariant elliptic cohomology classes on $T^*\mathcal{F}_\lambda$.

For $I \in \mathcal{I}_\lambda$ make the substitution

$$(7.6) \quad q, h, \mu_1/\mu_2, \dots, \mu_{N-1}/\mu_N \mapsto e^{2\pi i \tau}, e^{2\pi i y}, e^{2\pi i \nu_1}, \dots, e^{2\pi i \nu_{N-1}}, \\ t_1^{(k)}, \dots, t_{\lambda^{(k)}}^{(k)} \mapsto e^{2\pi i \nu_{1,1}}, \dots, e^{2\pi i \nu_{1,\lambda_1}}, e^{2\pi i \nu_{2,1}}, \dots, e^{2\pi i \nu_{2,\lambda_2}}, \dots, e^{2\pi i \nu_{k,1}}, \dots, e^{2\pi i \nu_{k,\lambda_k}},$$

for $k = 1, \dots, N$ in the function $\tilde{W}_{\sigma,I}^{\text{ell}}$. The resulting function is denoted by $\widehat{W}_{\sigma,I}^{\text{ell}}(\mathbf{v}, y, \boldsymbol{\nu})$. The function $\widehat{W}_{\sigma,I}^{\text{ell}}$ can be viewed as a section of a certain line bundle on $E^n \times E^{(\lambda_1)} \times \dots \times E^{(\lambda_N)} \times E \times E^{N-1}$. Its pull-back $c^*\widehat{W}_{\sigma,I}^{\text{ell}}$ by the embedding

$$c : \widehat{E}_T(T^*\mathcal{F}_\lambda) \rightarrow E^n \times E^{(\lambda_1)} \times E^{(\lambda_2)} \times \dots \times E^{(\lambda_N)} \times E \times E^{N-1}$$

is a section of the pull-back bundle and its restriction to $Y_J = \iota_J \widehat{E}_T(\text{pt})$ is the evaluation of $c^*\widehat{W}_{\sigma,I}^{\text{ell}}$ at $v^{(k)} = \mathbf{x}_{J_k}$, $k = 1, \dots, N$.

Recall the function $\psi_I(h, \boldsymbol{\mu})$ defined in (2.13). Make the substitution (7.6) in $\psi_I(h, \boldsymbol{\mu})$. Denote the resulting function by $\widehat{\psi}_I(y, \boldsymbol{\nu})$. The zeros of $\widehat{\psi}_I(y, \boldsymbol{\nu})$ define a divisor D_I on $\widehat{E}_T(\text{pt})$ consisting of hypertori with equation of the form $\nu_a + \nu_{a+1} + \dots + \nu_b - jy$, $i \in \mathbb{Z}$.

Let $M_I(\mathbf{x}, y, \boldsymbol{\nu})$ be the quadratic form defined in (6.4) and $\mathcal{L}_I = \mathcal{L}(M_I, 0)$ the corresponding line bundle on $\widehat{E}_T(\text{pt})$.

Proposition 7.1. *The restriction $c^*\widehat{W}_{\sigma,I}^{\text{ell}}$ of $\widehat{W}_{\sigma,I}^{\text{ell}}$ to $\widehat{E}_T(T^*\mathcal{F}_\lambda)$ is a meromorphic section of the admissible line bundle $p_T^*\mathcal{L}_{\sigma,I} \otimes \mathcal{T}_\lambda$ for some line bundle $\mathcal{L}_{\sigma,I}$ on $\widehat{E}(\text{pt})$. Moreover, if $\sigma = \text{id}$, the restriction $c^*\widehat{W}_{\text{id},I}^{\text{ell}} = c^*\widehat{W}_I^{\text{ell}}$ is a meromorphic section of the admissible line bundle $p_T^*\mathcal{L}_I \otimes \mathcal{T}_\lambda$, where \mathcal{L}_I is defined above, and the restriction is a holomorphic section of the line bundle $p_T^*\mathcal{L}_I(D_I) \otimes \mathcal{T}_\lambda$, where the notation $\mathcal{L}(D)$ means as usual the invertible sheaf of meromorphic sections of a sheaf \mathcal{L} whose poles are bounded by the divisor D .*

Proof. The proof is similar to the proof of [FRV, Proposition 5.9]. It is enough to prove the statement for $\sigma = \text{id}$.

The function $\widehat{G}_I(\mathbf{x}, y, \boldsymbol{\nu})\widehat{G}_\lambda(\mathbf{v}, y, \boldsymbol{\nu})/\widehat{E}_\lambda^{\text{ell}}(\mathbf{v}, y)$ defines a meromorphic section of the line bundle $\mathcal{L}(M_I + M_\lambda, 0)$ on $E^n \times E^{\lambda_1} \times \dots \times E^{\lambda_N} \times E \times E^{N-1}$. That line bundle lifts to the line bundle $p_T^*\mathcal{L}_I \otimes \mathcal{T}_\lambda$ on $\widehat{E}_T(T^*\mathcal{F}_\lambda)$. Lemmas 2.3 and 2.6 imply that $c^*\widehat{W}_I^{\text{ell}}$ is a meromorphic section of the admissible line bundle $p_T^*\mathcal{L}_I \otimes \mathcal{T}_\lambda$ on $\widehat{E}_T(T^*\mathcal{F}_\lambda)$. The fact that $c^*\widehat{W}_I^{\text{ell}}$ is a holomorphic section of the line bundle $p_T^*\mathcal{L}_I(D_I) \otimes \mathcal{T}_\lambda$ follows from the formula for the weight functions. \square

Corollary 7.2. *For any $\sigma \in S_n$ and $I \in \mathcal{I}_\lambda$ the restriction $c^*\widehat{W}_{\sigma,I}^{\text{ell}}$ defines a T -equivariant elliptic cohomology class on $T^*\mathcal{F}_\lambda$.*

7.7. Elliptic stable envelope for cotangent bundles of partial flag varieties. In this section we introduce a version of elliptic stable envelopes. Our elliptic stable envelopes are defined in terms of the elliptic weight functions. In Theorem 7.3 we give their axiomatic definition in the spirit of [MO1], [FRV]. *It would be interesting to understand the relation of our definition with the one sketched in [AO].*

For $\sigma \in S_N$, $I \in \mathcal{I}_\lambda$ we call the T -equivariant elliptic cohomology class $c^*\widehat{W}_{\sigma,I}^{\text{ell}}$ the *stable envelope associated with σ and the fixed point $x_I \in T^*\mathcal{F}_\lambda$* .

Consider \mathbb{C}^N with basis v_1, \dots, v_N . The standard basis of $(\mathbb{C}^N)^{\otimes n}$ is formed by the vectors v_I labeled by partitions $I = (I_1, \dots, I_N)$ of $\{1, \dots, n\}$, $v_I = v_{i_1} \otimes \dots \otimes v_{i_n}$, where $i_a = j$ if $a \in I_j$.

For $\text{id} \in S_n$ we may consider the map

$$(7.7) \quad \text{Stab}_{\text{id}} : (\mathbb{C}^N)^{\otimes n} \rightarrow \bigoplus_{\lambda \in (\mathbb{Z}_{\geq 0})^N, |\lambda|=n} \bigoplus_{I \in \mathcal{I}_\lambda} H_T^{\text{ell}}(T^*\mathcal{F}_\lambda)_{\mathcal{L}(D_I)},$$

sending v_I to the cohomology class $c^*\widehat{W}_I^{\text{ell}}$.

Similarly we may define the maps Stab_σ for all $\sigma \in S_n$. The maps $\text{Stab}_\sigma, \text{Stab}_{\sigma'}$ are related by the elliptic dynamical R -matrix, see Theorem 2.2, cf. [RTV2, Theorem 7.1] and Section 5.3.

Remark. For $\sigma \in S_n$ the collection $(c^*\widehat{W}_{\sigma,I}^{\text{ell}})_{I \in \mathcal{I}_\lambda}$ forms a basis of the T -equivariant cohomology of $T^*\mathcal{F}_\lambda$ in the sense of [FRV, Theorem 5.23]. We will discuss this fact in the next paper.

Remark. Using the map Stab_{id} one can construct an action of the dynamical elliptic quantum groups associated with \mathfrak{gl}_N on the extended equivariant cohomology

$\sqcup_{\lambda \in \mathbb{Z}_{\geq 0}^N, |\lambda|=n} \widehat{E}_T(T^*\mathcal{F}_\lambda)$. The action is by S_n -equivariant admissible difference operators acting on sections of admissible line bundles, see the case $N = 2$ in [FRV]. See similar constructions for equivariant cohomology and K -theory in [GRTV, RTV2, RV1]. We plan to discuss this action in the next paper.

Remark. The equivariant elliptic cohomology class $c^*\widehat{W}_I^{\text{ell}}$ has analogs in equivariant cohomology and equivariant K -theory of $T^*\mathcal{F}_\lambda$, see [GRTV, RTV1, RTV2]. The analog of $c^*\widehat{W}_I^{\text{ell}}$ in equivariant cohomology is the equivariant Chern–Schwartz–MacPherson class (or characteristic cycle) of the open Schubert variety Ω_I , see [RV2]. Hence $c^*\widehat{W}_I^{\text{ell}}$ may be viewed as an equivariant elliptic version of the Chern–Schwartz–MacPherson class.

7.8. Axiomatic definition of the stable envelope.

Theorem 7.3. *For any I the T -equivariant elliptic cohomology class $c^*\widetilde{W}_I^{\text{ell}}$ satisfies the following properties.*

- (i) *It is a meromorphic section of the admissible line bundle $p_T^*\mathcal{L}(N_I, 0) \otimes \mathcal{T}_\lambda$ for a suitable quadratic form N_I .*
- (ii) *The restriction of $c^*\widetilde{W}_I^{\text{ell}}$ to the component Y_I , written as a function $\mathbb{C}^{n+1+N-1} \rightarrow \mathbb{C}$ with transformation properties determined by the line bundle $p_T^*\mathcal{L}(N_I, 0) \otimes \mathcal{T}_\lambda$, equals*

$$(7.8) \quad \widehat{P}_I^{\text{ell}}(\mathbf{x}, y) = \prod_{k < l} \prod_{a \in I_k} \left(\prod_{\substack{b \in I_l \\ b < a}} \theta(y + x_b - x_a) \prod_{\substack{b \in I_l \\ b > a}} \theta(x_b - x_a) \right).$$

- (iii) *The restriction of $c^*\widetilde{W}_I^{\text{ell}}$ to any component Y_J , written as a function $\mathbb{C}^{n+1+N-1} \rightarrow \mathbb{C}$ with transformation properties determined by $p_T^*\mathcal{L}(N_I, 0) \otimes \mathcal{T}_\lambda$, is of the form*

$$(7.9) \quad \frac{1}{\widehat{\psi}_I(y, \boldsymbol{\nu})} \prod_{k < l} \prod_{a \in J_k} \prod_{\substack{b \in J_l \\ b < a}} \theta(y + x_b - x_a) \cdot F_{I,J},$$

where $F_{I,J}$ is a holomorphic function.

Moreover, these three properties uniquely determine the T -equivariant elliptic cohomology class $c^*\widetilde{W}_I^{\text{ell}}$.

Proof. The proof is analogous to the proof of [FRV, Theorem A.1].

Properties (i-iii) of $c^*\widetilde{W}_I^{\text{ell}}$ follow from Lemmas 2.5 and 2.6. Now we prove that properties (i-iii) uniquely determine $c^*\widetilde{W}_I^{\text{ell}}$. Let s be a section with properties (i-iii) for some admissible line bundle $p_T^*\mathcal{L}(N_I, 0) \otimes \mathcal{T}_\lambda$. Property (ii) implies that $N_I = M_I$, where M_I is given by (6.4). Hence $\mathcal{L}(N_I, 0) = \mathcal{L}_I$.

Denote by κ_I the difference between s and $c^*\widetilde{W}_I^{\text{ell}}$. Assume that the difference is nonzero. Then there exists a J such that κ_I restricted to Y_J is not 0. For a total ordering \prec refining the partial order $<$ on \mathcal{I}_λ let us choose J to be the largest index with the property $\kappa_I|_{Y_J} \neq 0$. We have $J \neq I$ because of the second property.

We claim that $\kappa_I|_{Y_J}$, written as a function $\mathbb{C}^{n+1+N-1} \rightarrow \mathbb{C}$, with transformation properties determined by $p_T^*\mathcal{L}_I \otimes \mathcal{T}_\lambda$, is of the form

$$(7.10) \quad \frac{1}{\widehat{\psi}_I(y, \nu)} \prod_{k < l} \prod_{a \in J_k} \left(\prod_{\substack{b \in J_l \\ b < a}} \theta(y + x_b - x_a) \prod_{\substack{b \in J_l \\ b > a}} \theta(x_b - x_a) \right) \cdot F_1,$$

where F_1 is holomorphic. The presence of the first theta-factors $\theta(y + x_b - x_a)$ in (7.10) follows from property (iii). The presence of the second theta-factors $\theta(x_b - x_a)$ in (7.10) follows from the fact that J is the largest index such that $\kappa_I|_{Y_J} \neq 0$, cf. the proof of [FRV, Theorem A.1].

Observe that the product of theta functions in (7.10) equals $c^*\widetilde{W}_J^{\text{ell}}|_J$ by property (ii). Hence $F_1/\widehat{\psi}_I$ is a meromorphic section of the line bundle $\mathcal{L}(M_I - M_J, 0)$.

Let $a \in \{1, \dots, n\}$ and $k, l \in \{1, \dots, N\}$, $k \neq l$, be such that $a \in I_k \cap J_l$. Consider $F_1/\widehat{\psi}_I$ as a function of x_a . Denote it by $f(x_a)$. The function $f(x_a)$ is a holomorphic function of x_a for generic fixed other arguments. Comparing the x_a -dependence of the quadratic form $M_I - M_J$ we obtain that

$$(7.11) \quad f(x_a + \tau) = e^{-2\pi i(2(\nu_l + \dots + \nu_{N-1} - \nu_k - \dots - \nu_{N-1}) + \dots)} f(x_a), \quad f(x_a + 1) = f(x_a),$$

where the dots indicate the terms independent on x_a and ν . Using the 1-periodicity, we expand $f(x_a) = \sum_{m \in \mathbb{Z}} a_m e^{2\pi i m x_a}$, and using the first transformation property of (7.11) we obtain that $a_m = 0$ for all $m \in \mathbb{Z}$. Hence $F_1 = 0$, and in turn, $\kappa_I|_{Y_J} = 0$. This is a contradiction proving that κ_I is 0 on all Y_J . \square

Remark. The classes appearing in the second and third properties in Theorem 7.3 can be identified as horizontal and vertical parts of the equivariant elliptic normal Euler classes of $C\Omega_{\sigma, I}$ near the fixed point x_I , cf. Section 4.6.

Remark. The stable envelopes are defined in equivariant cohomology, equivariant K -theory, and equivariant elliptic cohomology, see [MO1, MO2], for the cotangent bundles of partial flag varieties the definitions are discussed in [RTV1, RTV2, FRV] and in Sections 5, 7. In all three cases the definition consists of three axioms. One axiom says that the restriction $\kappa_J|_{x_J}$ of the stable envelope κ_J to the fixed point x_J equals some “expected” product. Another axiom says that the restriction $\kappa_J|_{x_I}$ of κ_J to any fixed point x_I should be divisible by some

product determined by the point x_I . This is a support type axiom, see Section 5. The last axiom says that the restriction $\kappa_J|_{x_I}$ should be “smaller” than the restriction $\kappa_I|_{x_I}$. The products in the first two axioms are the products of linear function or trigonometric functions or theta functions with the same arguments when we change from equivariant cohomology to equivariant K -theory and then to equivariant elliptic cohomology. It is interesting to see how the notion of “smallness” is changing in these three cases. In the cohomology case the restrictions are polynomials and one polynomial is smaller than another if the degree of the first is smaller than the degree of the second. In the K -theory case the restrictions are Laurent polynomials and one Laurent polynomial is smaller than another if the Newton polytope of the first can be parallelly moved inside the Newton polytope of the second. In the elliptic case the smallness is the requirement to be a section of an admissible line bundle, in other words, to be an equivariant elliptic cohomology class, see condition (i) in Theorem 7.3. In other words, the most nontrivial condition of smallness just dissolves in the definition of a cohomology class.

8. APPENDIX: COMPARISON OF BETHE ALGEBRAS

The Faddeev-Takhtajan-Reshetikhin formalism applied to the R -matrix $R(z, h)$ given by (5.5) produces the quantum loop algebra $\mathcal{U}(\widetilde{\mathfrak{gl}}_N)$, the evaluation N -dimensional representations of $\mathcal{U}(\widetilde{\mathfrak{gl}}_N)$, and commutative Bethe subalgebras \mathcal{B}^q of $\mathcal{U}(\widetilde{\mathfrak{gl}}_N)$ depending on complex parameters $q = (q_1, \dots, q_N)$ called the quantum parameters, see explicitly these constructions in [RTV2, Sections 10 and 11]. The Bethe algebra acts on the tensor product of n evaluation representations and preserves the weight decomposition $(\mathbb{C}^N)^{\otimes n} = \bigoplus_{\lambda \in \mathbb{Z}_{\geq 0}^N, |\lambda|=n} (\mathbb{C}^N)_{\lambda}^{\otimes n}$. The image of the Bethe algebra in the algebra of endomorphisms of a weight subspace $(\mathbb{C}^N)_{\lambda}^{\otimes n}$ is described in [RTV2, Theorem 13.3] by generators and relations in terms of a discrete Wronski map depending on parameters q .

Let $\mathcal{R}(z, h)$ be defined by (5.4) for the anti-dominant alcove Δ , i.e. with all $m_{j,k}$ equal to zero. The matrix $\mathcal{R}(z, h)$ differs from the matrix $R(z, h)$ by change in the normalization convention involving h . The Faddeev-Takhtajan-Reshetikhin formalism applied to $\mathcal{R}(z, h)$ produces the new quantum loop algebra $\mathcal{U}(\widetilde{\mathfrak{gl}}_N)'$ isomorphic to $\mathcal{U}(\widetilde{\mathfrak{gl}}_N)$, new evaluation representations, and new commutative Bethe subalgebras $\mathcal{B}^{q'} \subset \mathcal{U}(\widetilde{\mathfrak{gl}}_N)'$. It turns out that *the image of the new Bethe algebra $\mathcal{B}^{q'}$ with parameters $q' = (q'_1, \dots, q'_N)$ in the algebra of endomorphisms of $(\mathbb{C}^N)_{\lambda}^{\otimes n}$ coincides with the image of the original Bethe algebra \mathcal{B}^q with quantum parameters $q = (q_1, \dots, q_N)$ if*

$$q'_k = h^{-\lambda_1 - \dots - \lambda_{k-1} + \lambda_{k+1} + \dots + \lambda_N} q_k, \quad k = 1, \dots, N.$$

Hence the image of the new Bethe algebra is also described by the discrete Wronski map. We will discuss this fact in detail in the next paper.

The importance of that statement lies in the following. In [OS] Okounkov and Smirnov consider the equivariant K -theory algebra of a Nakajima variety and the stable envelopes of the anti-dominant alcove. Using the R -matrices of that alcove they define the associated quantum loop algebra with the action on the equivariant K -theory algebra. Conjecturally, the associated Bethe algebra of that action is the algebra of quantum multiplication on the equivariant K -theory algebra, cf. [OS, RTV2, PSZ]. For the cotangent bundle $T^*\mathcal{F}_{\lambda}$ of a partial flag variety \mathcal{F}_{λ} the constructions of this paper and [OS, RTV2] identify the Bethe

algebra of the cotangent bundle with the image of the Bethe algebra \mathcal{B}'^d in the endomorphism algebra of $(\mathbb{C}^N)_\lambda^{\otimes n}$. Hence the formulated statement gives a conjectural description of the algebra of quantum multiplication on the equivariant K -theory algebra of $T^*\mathcal{F}_\lambda$ by generators and relations in terms of the discrete Wronski map.

REFERENCES

- [AO] M. Aganagic, A. Okounkov, *Elliptic stable envelopes*, Preprint 2016, arXiv:1604.00423
- [ChG] N. Chriss, V. Ginzburg, *Representation theory and complex geometry*, Modern Birkhäuser Classics, Birkhäuser Inc., Boston, MA, 2010
- [F] G. Felder, *Elliptic quantum groups*, in Proceedings of the ICMP, Paris 1994 (D. Iagolnitzer, ed.), Intern. Press, Cambridge, MA, 1995 pp. 211–218
- [FR] L. Feher, R. Rimanyi, *Calculation of Thom polynomials and other cohomological obstructions for group actions*, Real and Complex Singularities (Sao Carlos, 2002), Ed. T. Gaffney and M. Ruas, Contemp. Math., #354, Amer. Math. Soc., Providence, RI, June 2004, pp. 69–93.
- [FRV] G. Felder, R. Rimanyi, A. Varchenko, *Elliptic dynamical quantum groups and equivariant elliptic cohomology*, Preprint 2017, arXiv:1702.08060
- [FTV1] G. Felder, V. Tarasov, A. Varchenko, *Solutions of the elliptic QKZB equations and Bethe ansatz I*, in Topics in Singularity Theory, V.I. Arnold’s 60th Anniversary Collection, Advances in the Mathematical Sciences, AMS Translations, Series 2, v.180, 45–76, 1997
- [FTV2] G. Felder, V. Tarasov, and A. Varchenko, *Monodromy of solutions of the elliptic quantum Knizhnik-Zamolodchikov-Bernard difference equations*, Int. J. Math. 10 (1999) 943–975
- [FP] W. Fulton, P. Pragacz, *Schubert Varieties and Degeneracy Loci, Appendix J by the authors in collaboration with I. Ciocan-Fontanine*, Lect. Notes in Math **1689** (1998), 1–148
- [Ga] Nora Ganter, *The elliptic Weyl character formula*, Compos. Math. 150 (2014), no. 7, 1196–1234
- [GKV] V. Ginzburg, Michail Kapranov, and Eric Vasserot, *Elliptic Algebras and Equivariant Elliptic Cohomology I*, arXiv:q-alg/9505012
- [GRTV] V. Gorbounov, R. Rimányi, V. Tarasov, A. Varchenko, *Cohomology of the cotangent bundle of a flag variety as a Yangian Bethe algebra* J. Geom. Phys. **74** (2013), 56–86
- [GKM] Mark Goresky, Robert Kottwitz, and Robert MacPherson, *Equivariant cohomology, Koszul duality, and the localization theorem*, Invent. Math., 131(1)(1998), 25–83
- [Gr] I. Grojnowski, *Delocalized equivariant elliptic cohomology*, (preprint 1994), in *Elliptic cohomology*, 114–121, London Math. Soc. Lecture Note Ser., 342, Cambridge Univ. Press, Cambridge, 2007.
- [MO1] D. Maulik, A. Okounkov, *Quantum Groups and Quantum Cohomology*, Preprint (2012), 1–276, arXiv:1211.1287
- [MO2] D. Maulik and A. Okounkov: in preparation; 2015
- [MTV] E. Mukhin, V. Tarasov, A. Varchenko, *Bethe eigenvectors of higher transfer matrices*, J. Stat. Mech. (2006), no. 8, P08002, 1–44

- [Mu] David Mumford, *Abelian varieties*, Tata Institute of Fundamental Research Studies in Mathematics, No. 5, Oxford University Press, London 1970
- [O] A. Okounkov. Lectures on K -theoretic computations in enumerative geometry. ArXiv: 1512.07363.
- [OS] Andrei Okounkov, Andrey Smirnov. Quantum difference equation for Nakajima varieties, Preprint 2016, arXiv:1602.09007
- [PSZ] Petr P. Pushkar, Andrey Smirnov, Anton M. Zeitlin, *Baxter Q -operator from quantum K -theory*, Preprint 2016, arXiv:1612.08723
- [RTV1] R. Rimányi, V. Tarasov, A. Varchenko, *Partial flag varieties, stable envelopes and weight functions*, Preprint (2012), 1–22; arXiv:1212.6240, Quantum Topol. 6 (2015), no. 2, 333–364, DOI 10.4171/QT/64
- [RTV2] R. Rimányi, V. Tarasov, A. Varchenko, *Trigonometric weight functions as K -theoretic stable envelope maps for the cotangent bundle of a flag variety*, arXiv:1411.0478, Journal of Geometry and Physics, v. **94** (2015), 81–119, doi:10.1016/j.geomphys.2015.04.002
- [RV1] R. Rimányi, A. Varchenko, *Dynamical Gelfand-Zetlin algebra and equivariant cohomology of Grassmannians*, arXiv:1510.03625, Journal of Knot Theory and Its Ramifications Vol. **25**, No. 12 (2016) (29 pages), DOI: 10.1142/S021821651642013X
- [RV2] R. Rimányi, A. Varchenko, *Equivariant Chern-Schwartz-MacPherson classes in partial flag varieties: interpolation and formulae*, arXiv:1509.09315, to appear in IM-PANGA2015, 9pp
- [RoKu] I. Rosu, *Equivariant K -theory and equivariant cohomology, with an appendix by Allen Knutson and Ioanid Rosu*, Math. Z. **243** (2003), no. 3, 423–448
- [SV] V. Schechtman and A. Varchenko, *Arrangements of Hyperplanes and Lie Algebra Homology*, Invent. Math. Vol. 106 (1991), 139–194
- [TV1] A. Varchenko, V. Tarasov, *Jackson integral representations for solutions of the Knizhnik-Zamolodchikov quantum equation*, Leningrad Math. J. **6** (1994), 275–313
- [TV2] V. Tarasov and A. Varchenko, *Geometry of q -hypergeometric functions as a bridge between Yangians and quantum affine algebras*, Invent. Math. **128** (1997), no. 3, 501–588
- [TV3] V. Tarasov and A. Varchenko, *Geometry of q -hypergeometric functions, quantum affine algebras and elliptic quantum groups*, Asterisque, **246** (1997), 1–135
- [TV4] V. Tarasov, A. Varchenko, *Combinatorial formulae for nested Bethe vectors*, SIGMA **9** (2013), 048, 1–28
- [V] A. Varchenko, *Quantized KZ equations, Quantum YBE, and difference equations for q -Hypergeometric functions*, Comm. Math. Phys. **162** (1994), 499–528